

EDUCATIONAL SOCIAL SCIENCE DEVELOPMENT PLANNING

Jurnal Scientia




	Register	Login
	ISSN : 2302-0059 (Print)	
https://infor.seaninstitute.org/index.php/pendidikan	Published by Sean Institute editorjurnal@seaninstitute.org	
Current Archives About Policies 🗸	About - Contac	t
S	earch	

Home / Editorial Team

Editorial Team

Editor in Chief






Associated Editorial











4





Sanjaya Pinem,S,Kom M.Cs


Scopus ID



Scholar ID

National Chung Cheng University





. Scopus ID

. Scholar ID

. -

. -

. Stmik Pelita

Nusantara,Indonesia

Stmik Pelita Nusantara,Indonesia

Amran Sitohang, . M.Kom

Additional Menu

Editor Team

Peer Review Process

Riviewer

Focus And Scope

Author Guidelines

Online Submission

Pulication Ethics

Journal Fee

CopyRight Notice

Statitistic View

Additional iv



ISSN:

P-ISSN: 2302-0059 E-ISSN: 2723-7486

TEMPLATE





SERTIFIKAT

Kementerian Riset dan Teknologi/ Badan Riset dan Inovasi Nasional



Petikan dari Keputusan Menteri Riset dan Teknologi/ Kepala Badan Riset dan Inovasi Nasional Nomor 200/M/KPT/2020 Peringkat Akreditasi Jurnal Ilmiah Periode III Tahun 2020 Nama Jurnal Ilmiah

Jurnal Scientia

E-ISSN: 27237486

Penerbit: SEAN INSTITUTE

Ditetapkan sebagai Jurnal Ilmiah

TERAKREDITASI PERINGKAT 5


Akreditasi Berlaku selama 5 (lima) Tahun, yaitu Volume 7 Nomor 2 Tahun 2019 sampai Volume 12 Nomor 1 Tahun 2024

Jakarta, 23 Desember 2020 Menteri Riset dan Teknologi/ Vepsta Bodian Piset dan Inovasi Nasional Republik Indonesia, Mutur

Bambang P. S. Brodjonegoro



Language





Jurnal Scientia

Sean Institute

Alamat: Komplek New Pratama ASri Blok C, No.2, Deliserdang, Sumatera Utara, Indonesia email : scientia@seaninstitute.org



Jurnal Scientia

is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Policies and Regulations Link

- Advertising Policy
- Corrections, Retractions & Expressions of Concern
- Editorial & Peer Review Process Stats
- Editorial Policies
- Data Sharing Solicy
- Informed Consent policy
- License Information
- Peer Review Policy
- Plagiarism Policy
- Privacy Policy



View My

- Publishing Ethics
- Publishing with Iocspublisher: step-by-step
- Policy of Screening for Plagiarism
- Policies on Conflict of Interest, Human and Animal rights, and Informed Consent
- Self-Archiving Policies
- Statement of Informed Consent
- Open Access

Download Pdf

: https://doi.org/10.58471/scientia.v12i03.1696 BAbstract views: 10 , BDownload Pdf downloads: 2

DETECT AND MITIGATE MALWARE THREATS USING SANDBOXING TECHNOLOGY

Ade Andryani, Tata Sutabri 3263-3269

Download Pdf

: https://doi.org/10.58471/scientia.v12i03.1689

Babstract views: 2 , BDownload Pdf downloads: 1

REVIEW OF POLYGRAPH INSTRUMENTS AS EVIDENCE IN COURT IN FULFILLING THE CRIMINAL PROCEDURE CODE JO LAW NUMBER 19 OF 2016 CONCERNING AMENDMENTS TO LAW NUMBER 11 OF 2008 CONCERNING ELECTRONIC INFORMATION AND TRANSACTIONS

Berti Nora Panjaitan, Chitto Chumbradika

3270-3276

Download Pdf

: https://doi.org/10.58471/scientia.v12i03.1690 ■Abstract views: 1 , Download Pdf downloads: 1

Animated Film " Lorong Waktu Si AA: Rafathar Cinta Batik Indonesia" As a Media to Instill a Sense of Love for Culture To Children

Kelvin Alfi Nurdiansyah

Download Pdf

: https://doi.org/10.58471/scientia.v12i03.1691 ■Abstract views: 0 , D Download Pdf downloads: 1

ANALYSIS OF THE CAUSES OF DEFECTS IN THE PRODUCTION PROCESS OF PILLOWCASES AND BOLSTERS USING THE FMEA METHOD APPROACH AT UD ARJUNO

Fatkhul Musif, Akhmad Wasiur Rizqi, Elly Ismiyah 3280-3288

Download Pdf

: https://doi.org/10.58471/scientia.v12i03.1692 ≧Abstract views: 0 , ⊵Download Pdf downloads: 1

Development of Amplification Refractory Mutation System PCR to Detect Androgen Receptor Gene G1773A (rs6152) Polymorphism Triyana Sari, Erick Sidarta, Sari Mariyati Dewi Nataprawira, Giovani Evangelista Alfonsius, Damar Sajiwo

Download Pdf

: https://doi.org/10.58471/scientia.v12i03.1694 ■Abstract views: 0 , D Download Pdf downloads: 1

CAPITAL STRUCTURE AND FIRM PERFORMANCE: AN EVIDENCE OF TRANSPORTATION, LOGISTIC AND WAREHOUSE COMPANY LISTED AT IDX 2018-2022

Ayu Shafira Rachmani, Eko Rizkianto

Download Pdf

: https://doi.org/10.58471/scientia.v12i03.1703 ÈAbstract views: 1 , ⊵Download Pdf downloads: 1

ANALYSIS OF THE EFFECT OF EXPORTS, IMPORTS, AND INFLATION RATE ON FOREIGN EXCHANGE RESERVES IN INDONESIA

Abidah Dwi Rahmi Satiti, Rivatul Ridho Elvierayani, Sukmawati Indah Dzikria

Download Pdf

BAbstract views: 0 , BDownload Pdf downloads: 2

LEGAL PROTECTION FOR THE FAMILIES OF TERRORISM CRIMINAL SUSPECTS

Yeni Nuraeni, Lasmin Alfies Sihombing

🕒 Download Pdf

Abstract views: 1 , BDownload Pdf downloads: 2

ASSISTANCE FOR COMMUNITY GROUP THROUGH ENVIRONMENTAL HEALTH EDUCATION IN SANCA VILLAGE, CIATER DISTRICT, SUBANG REGENCY

Riany L Nurwulan, Abu Huraerah

₿ 3319-3324

Download Pdf

Abstract views: 0 , Bownload Pdf downloads: 2

http://infor.seaninstitute.org/index.php JURNAL SCIENTIA, Volume 12 No 03, 2023

DEVELOPMENT OF AMPLIFICATION REFRACTORY MUTATION SYSTEM PCR TO DETECT ANDROGEN RECEPTOR GENE G1773A (RS6152) POLYMORPHISM

Triyana Sari¹, Erick Sidarta², Sari Mariyati Dewi Nataprawir³, Giovani Evangelista Alfonsius⁴, Damar Sajiwo⁵

^{1,2,3,4}Universitas Tarumanagara, Jakarta, Indonesia, ⁵NMW Clinic, Jakarta, Indonesia Email: triyanas@fk.untar.ac.id

Article Info

Received: 22/06/2023 Revised: 19/07/2023 Accepted: 24/07/2023 Androgen receptor (AR) is a ubiquitous receptor responsible for responses by androgen stimulus. Androgen, a hormone which will bind to the AR, is essentials for normal male sexual development. Nevertheless, one of the polymorphisms in the AR gene, G1733A (rs6152) have been associated with numerous clinical risks such as cardiovascular disease (CD), androgenetic alopecia, high prostate-specific antigen (PSA) levels, male infertility, recurrent spontaneous abortions and prostate cancer. This study aims to develop an alternative and cost-effective method to detect G1773A (rs6152) polymorphism. In this study, amplification refractory mutation system-polymerase chain reaction (ARMS-PCR) using two pair of primers was used to detect the G1733A (rs6152) polymorphism. Primer design was done using http://primer1.soton.ac.uk/primer1.html online tools and then manually adjusted to increase the specificity. A total of 54 samples were screened using ARMS PCR and 2 representative samples of each allele from previous screening were used to validate the results using Sanger DNA sequencing. Among 54 subjects screened, we found 52 (96.3%) subjects carry G allele and 2 (3.7%) subjects carry A allele. No heterozygote was found in this study. The frequency of G allele was 96.97% and the frequency of A allele was 3.03%. Result validation using DNA sequencing was in agreement with ARMS-PCR method results. ARMS-PCR can be used as efficient alternative for genotyping of G1733A (rs6152) AR gene polymorphism.

Keywords: Androgen receptor, rs6152, ARMS-PCR, polymorphism

1. Introduction

Androgen, such as testosterone and dihydrotestosterone (DHT), were steroid hormones involved in the development of male reproductive organs, such as: embryogenesis; puberty and sexual maturation.(1) In addition, androgen also have diverse effect in bone and skeletal muscle, either in male or female.(2) The mechanisms of androgen action involves the enzymatic process of reduction of testosterone by 5_{α} -reductase to DHT and followed by activation of Androgen Response Elements (ARE), a specific motif presents in promoter or enhancer which activation rely on androgen receptor (AR).(3) Nevertheless, the binding of testosterone, DHT and activation of ARE were mediated by AR.

The AR is a receptor belong to superfamily of DNA binding transcription factors which share both DNA sequence and functional homology such as steroid, thyroid, vitamin D, retinoic acid hormones and many others. AR gene is located on the X chromosome on Xq11- 12. Significant effect due to the mutations in the AR may result in androgen insensitivity syndrome (AIS) which characterized by evidence of feminization of the male genitals at birth, abnormal secondary sexual development during puberty and infertility in individuals with normal 46XY chromosomes. The clinical presentation of AIS is vary with different phenotype due to large amount of mutation and heterogeneity (missense mutations, nonsense mutations, splicing variant, deletions and insertions) in the AR gene.(4) One of the mutations in AR, G1733A (rs6152) have been associated with increased risks of many clinical outcomes such as androgenetic alopecia (AGA),(5) higher prostate specific antigen (PSA),(5) recurrent spontaneous abortion,(6) polycystic ovary syndrome,(7) cardiovascular disease(8) and prostate cancer aggressiveness.(9)

🕞 🛈 😌 Jurnal Scientia is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-



http://infor.seaninstitute.org/index.php

JURNAL SCIENTIA, Volume 12 No 03, 2023

Currently, the most frequent method to detect G1733A (rs6152) polymorphism is by using polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) with the help of *StuI* restriction enzyme(5,8). This method involves two different process in which amplification of target DNA sequence and followed by cutting of the amplified band using specific enzyme. Here we introduce Amplification Refractory Mutation System (ARMS) which is a reliable, simple and fast method to detect the SNPs by using four primers in one PCR reaction.

2. Method

Samples and DNA extraction

A total of 54 subjects were used in this study. Informed consent was given to each subjects and ethical clearance were given by Universitas Tarumanagara Human Research Ethics Committee with ethical clearance number PPZ20222090. Blood DNA extraction were done using Genomic DNA Midi Kit (Geneaid, Taiwan). After extraction, DNA quantification were done by spectrophotometer (Thermo Fischer Scientific), each sample then diluted to approximately 50 ng/mL DNA concentration and kept at -20°C until further use.

Tetra arms PCR for rs6152 Four primers were designed by PRIMER1 online software (<u>http://primer1.soton.ac.uk/primer1.html</u>). Further customization was done manually using Bioedit software ver 7.2.5 (<u>https://bioedit.software.informer.com/7.2/</u>) to increase the specificity of the primers (Table 1). ARMS-PCR were done in total volume of 25 μ L with 12.5 μ L 2x myTaq TM mastermix (Meridian Bioscience, United Kingdom), 0.4 μ M of FO primer, 0.4 μ M of RO primer, 0.4 μ M of FI primer, 0.4 μ M of RI primer and topped with dH2O. The PCR conditions are initial denaturation at 95°C for 2 minutes, 30 cycles of denaturation at 95°C for 30 seconds, annealing at 63°C for 30 seconds, extension at 72°C for 10 seconds and followed by final extention at 72°C for 5 minutes at the end of cycle. The PCR products were visualized by 1.5% agarose gel electrophoresis at 100 V for 45 minutes.

Primer	Sequence (5'- 3')	Product size
Forward inner (FI)	CAGCAGCGGGAGAGCGAGGTAG	G allele 245 bp
Reverse inner (RI)	AAGTGGGAGCCCCCGAGTCT	A allele 367 bp
Forward outer (FO)	CAAGCCCATCGTAGAGGCCC	Outer band 572 bp
Reverse outer (RO)	GCCAATGGGGCACAAGGAGT	_

DNA Sequencing validation

DNA Sequencing method was employed to validate the result of ARMS-PCR. Four samples were chosen randomly from the G allele positive ARMS-PCR and all samples (n = 2) from A allele positive ARMS-PCR, which is the minor allele, were amplified using only the outer primer (FO & RO) and then outsourced for the sequencing process (Apical Scientific Laboratory, Malaysia).



http://infor.seaninstitute.org/index.php JURNAL SCIENTIA, Volume 12 No 03, 2023

3. Result And Discussion




Figure 1. Electropherogram of rs6152 ARMS-PCR. lane M is for DNA ladder, lane 1, 3 and 5 are products of the G allele positive with two bands at 572 and 245 bp, lane 2 and 4 were products of the A allele positive with three bands at 572 bp, 375 bp and one unspecific band, lane 6 was negative controls.

A representative for the rs6152 ARMS-PCR were shown in figure 1. The G allele shows two bands which are 572 and 245 bp, while A allele shows two band with different pattern which are 572 and 367 bp. Among 54 samples enrolled in this study, 52 (96.3%) were G allele positive and 2 (3.7%) were A allele positive. We cannot categorize the allele assignment by genotype (GG/GA/AA) due to rs6152 was sex linked SNPs. No heterozygotes in the female subjects were found in this study (table 1). The allele frequencies were 96.97% for the G allele and 3.03% for the A allele. The distribution of the genotypes in this study was not in Hardy-Weinberg equilibrium ($\chi^2 = 7.68287$; P = 0.021)

Table 2. Allele and genotype distribution in this study		
Genotype	Distribution	
Female (n=45)		
GG	44 (97.78%)	
AA	1 (2.22%)	
Male (n=9)		
G	8 (88.89%)	
А	1 (11.11%)	
Allele Frequency		
G	96 (96.97%)	
Α	3 (3.03%)	

The representative results of DNA sequencing are shown in Figure 2. Two different samples with different result, the G allele positive (572 and 245 bp) were randomly chosen and both samples with A allele positive (572 and 375 bp), were used for validation by sequencing. The results of ARMS PCR agreed with the DNA sequencing results.




Figure 2. DNA sequencing result of rs6152. A, G allele positive; B, A allele positive

Discussion

Androgen receptor, which gene is located on X chromosome Xq11-12, is a receptor that stimulated by testosterone and 5_{α} -DHT. Both hormone function as a ligand dependent nuclear transcription factor which affect the activation for the AR. AR is expressed widely in many cells and tissues. it modulates diverse range of biological mechanisms, including the development and maintenance of reproductive, cardiovascular, neural, musculoskeletal, immune and hemopoetic system.(10) Genome-wide association studies (GWAS) have identified a number of SNPs in this gene and its possible associations with higher risks of clinical outcomes such as prostate cancer,(11) AIS,(4) autism,(12) polycystic ovary syndrome (7) and many more. One of the SNP located in the AR gene, rs6152 is one of the mutation that have been associated with various clinical outcomes.(5–8)

ARMS PCR is already reported as a simple, fast and low-cost method in SNP genotype detection and this method already been compared to PCR-RFLP, which need restriction enzyme and a two process method. The application of ARMS PCR is already widely used in detecting various SNPs.(13–18) This method uses four primer in single PCR reaction in which a pair of the primers recognize one specific SNP.(19) The challenge in this method was the primer design and the optimization process. The primers used in this study were designed by PRIMER1 software. After the *insilico* primer design was recommended by the software, we use BLAST to see the specificity of the primers. The inner primer which was supposed to detect the G-allele shown the specificity for the *homo sapiens* chromosome 17. To overcome this issue, we altered the inner G-allele primer manually by adding 2 nucleotides to increase the specificity for the AR gene.

The problem following the issues is the high GC content of each primers (ranged from 60-70%) due to the high GC of rs6152 site, however the Tm result from the high GC content didn't exceed 67. Nevertheless, to ensure a working PCR reaction, we choose a pre-mixed PCR mastermix which can handle high GC content reaction and known for its capability in multiplex PCR. ARMS PCR were not recommended for detection high GC percentage region by Medrano and Oliveira, (19) but recent study shown the use of dimethyl sulfoxide as PCR enhancer or thermostable strand displacement polymerase can overcome the problem.(14) As shown in figure 1, The G allele (245 bp) and A allele (367 bp) shown the same intensity of band luminance which shown the PCR reaction is working. Subjects number 2 and 4 also shown an unspecific band with lower luminance and higher placement than the G allele band which can be ignored. Unspecific bands were common in ARMS PCR and various method may be employed to eliminate the unspecific band.(19) Although no heterozygote alleles were found in this study, this method also capable to detect heterozygote allele (GA) which was obtained by mixing 2 different samples with different alleles (data not shown). Lastly, to ensure the validity of the ARMS PCR for rs6152, we sequenced 4 random samples from samples shown ARMS PCR G allele positive and 2 samples of the ARMS PCR A allele positive in which the sequencing results were in concordant with the ARMS PCR results.

According to our knowledge, this is the second time the detection of rs6152 were done in Indonesia and the first study of rs6152 SNP prevalence for healthy people. The only known study for rs6152 were done by Sribudiani et al. (2018) which detect the prevalence of rs6152 SNP in prostate cancer patients. Our study shows lower minor allele frequency (MAF) of rs6152 (0.03) than the study reported by Sribudiani et al. (2018) in prostate cancer patients (0.0789).(20)

🕞 🛈 😔 Jurnal Scientia is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-

http://infor.seaninstitute.org/index.php

JURNAL SCIENTIA, Volume 12 No 03, 2023

4. Conclusion

ARMS-PCR, is an effective, simple, reliable and low cost SNP detection method for rs6152 despite its high GC content site, which can be used as effective screening method as an alternative to PCR-RFLP. This study also shown low MAF (A allele) of rs6152 in Indonesian population. We would like to give our gratitude to dr. Nataliani Mawardi and NMW Clinics for the help in providing subjects for this study. We also want to give our gratitude to Universitas Tarumanagara for funding this research and Laboratorium Terpadu FK Untar for the laboratory use.

References

- 1. Murashima A, Kishigami S, Thomson A, Yamada G. Androgens and mammalian male reproductive tract development. Biochim Biophys Acta Gene Regul Mech. 2015 Feb 1;1849(2):163–70.
- 2. Notelovitz M. Androgen effects on bone and muscle. Fertil Steril [Internet]. 2002 [cited 2022 Nov 28];77(SUPPL. 4):34–41. Available from: https://pubmed.ncbi.nlm.nih.gov/12007900/
- 3. Wilson S, Qi J, Filipp F V. Refinement of the androgen response element based on ChIP-Seq in androgen-insensitive and androgen-responsive prostate cancer cell lines. Sci Rep [Internet]. 2016 Sep 14 [cited 2022 Nov 28];6. Available from: https://pubmed.ncbi.nlm.nih.gov/27623747/
- 4. Gulía C, Baldassarra S, Zangari A, Briganti V, Gigli S, Gaffi M. Androgen insensitivity syndrome.
- 5. Kucerova R, Bienova M, Kral M, Bouchal J, Trtkova KS, Burdova A, et al. Androgenetic alopecia and polymorphism of the androgen receptor gene (SNP rs6152) in patients with benign prostate hyperplasia or prostate cancer. J Eur Acad Dermatol Venereol [Internet]. 2015 Jan 1 [cited 2022 Oct 13];29(1):91–6. Available from: https://pubmed.ncbi.nlm.nih.gov/24665929/
- Porras-Dorantes Á, Brambila-Tapia AJL, Lazcano-Castellanos AB, Da Silva-José TD, Juárez-Osuna JA, García-Ortiz JE. Association between G1733A (rs6152) polymorphism in androgen receptor gene and recurrent spontaneous abortions in Mexican population. J Assist Reprod Genet [Internet]. 2017 Oct 1 [cited 2022 Feb 26];34(10):1303. Available from: /pmc/articles/PMC5633579/
- Peng CY, Long XY, Lu GX. Association of AR rs6152G/A gene polymorphism with susceptibility to polycystic ovary syndrome in Chinese women. Reprod Fertil Dev [Internet]. 2010 May 20 [cited 2022 Nov 29];22(5):881–5. Available from: https://www.publish.csiro.au/rd/RD09190
- 8. Agiannitopoulos K, Bakalgianni A, Marouli E, Zormpa I, Manginas A, Papamenzelopoulos S, et al. Gender Specificity of a Genetic Variant of Androgen Receptor and Risk of Coronary Artery Disease. J Clin Lab Anal [Internet]. 2016 May 1 [cited 2022 Feb 26];30(3):204. Available from: /pmc/articles/PMC6807026/
- Robles-Fernandez I, Martinez-Gonzalez LJ, Pascual-Geler M, Cozar JM, Puche-Sanz I, Serrano MJ, et al. Association between polymorphisms in sex hormones synthesis and metabolism and prostate cancer aggressiveness. PLoS One [Internet]. 2017 Oct 1 [cited 2022 Nov 29];12(10):e0185447. Available from:

https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0185447

- Rana K, Davey RA, Zajac JD. Human androgen deficiency: insights gained from androgen receptor knockout mouse models. Asian J Androl [Internet]. 2014 Mar [cited 2022 Feb 26];16(2):169–77. Available from: https://pubmed.ncbi.nlm.nih.gov/24480924/
- 11. Morova T, McNeill DR, Lallous N, Gönen M, Dalal K, Wilson DM, et al. Androgen receptorbinding sites are highly mutated in prostate cancer. Nat Commun [Internet]. 2020 Dec 1 [cited 2022 Dec 20];11(1). Available from: /pmc/articles/PMC7012874/
- 12. Henningsson S, Jonsson L, Ljunggren E, Westberg L, Gillberg C, Råstam M, et al. Possible association between the androgen receptor gene and autism spectrum disorder.

Jurnal Scientia is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-



http://infor.seaninstitute.org/index.php

JURNAL SCIENTIA, Volume 12 No 03, 2023

Psychoneuroendocrinology [Internet]. 2009 Jun [cited 2022 Dec 20];34(5):752–61. Available from: https://pubmed.ncbi.nlm.nih.gov/19167832/

- Dewi AL, Paramita DK, Facliiroli J. Development of Tetra-primer Amplification Refractory Mutation System (ARMS) PCR for Detection of CHRNA3 rs8040868. Indones Biomed J [Internet]. 2021 Jun 14 [cited 2022 Dec 20];13(2):192–200. Available from: https://inabj.org/index.php/ibj/article/view/1463
- 14. Alyethodi RR, Singh U, Kumar S, Alex R, Deb R, Sengar GS, et al. T-ARMS PCR genotyping of SNP rs445709131 using thermostable strand displacement polymerase. BMC Res Notes [Internet]. 2018 Feb 15 [cited 2022 Dec 20];11(1):1–5. Available from: https://bmcresnotes.biomedcentral.com/articles/10.1186/s13104-018-3236-6
- 15. Mehdi M, 1□ H, Hadadzadeh M, Fallahzadeh H. Development of One-Step Tetra-primer ARMS-PCR for Simultaneous Detection of the Angiotensin Converting Enzyme (ACE) I/D and rs4343 Gene Polymorphisms and the Correlation with CAD Patients. Avicenna J Med Biotechnol [Internet]. 2019 [cited 2022 Dec 20];11(1):118. Available from: /pmc/articles/PMC6359703/
- Jin C, Li Z, Zheng X, Shen K, Chao J, Dong Y, et al. Development and validation of T-ARMS-PCR to detect CYP2C19*17 allele. J Clin Lab Anal [Internet]. 2020 Jan 1 [cited 2022 Dec 20];34(1). Available from: /pmc/articles/PMC6977150/
- 17. Yang L, Ijaz I, Cheng J, Wei C, Tan X, Khan MA, et al. Evaluation of amplification refractory mutation system (ARMS) technique for quick and accurate prenatal gene diagnosis of CHM variant in choroideremia. Appl Clin Genet [Internet]. 2017 Dec 19 [cited 2023 Jan 16];11:1–8. Available from: https://www.dovepress.com/evaluation-of-amplification-refractory-mutation-system-arms-technique--peer-reviewed-fulltext-article-TACG
- Garbuz MM, Ovchinnikova AA, Kumeiko VV. Design, Optimization and Validation of the ARMS PCR Protocol for the Rapid Diagnosis of Wilson's Disease Using a Panel of 14 Common Mutations for the European Population. Genes 2022, Vol 13, Page 1940 [Internet]. 2022 Oct 25 [cited 2023 Jan 16];13(11):1940. Available from: https://www.mdpi.com/2073-4425/13/11/1940/htm
- Medrano RFV, De Oliveira CA. Guidelines for the Tetra-Primer ARMS–PCR Technique Development. Mol Biotechnol 2014 567 [Internet]. 2014 Feb 12 [cited 2022 Dec 20];56(7):599–608. Available from: https://link.springer.com/article/10.1007/s12033-014-9734-4
- 20. Sribudiani Y, Marwan DW, Aulanni'am A, Widodo MA, Purnomo BB, Panigoro R, et al. Germline Mutations and Polymorphisms of Androgen Receptor in Prostate Cancer Patients: Frequency and Results of in Silico Analysis. Asian Pac J Cancer Prev [Internet]. 2018 Aug 1 [cited 2022 Dec 20];19(8):2241. Available from: /pmc/articles/PMC6171394/