



PROSIDING

SEMINAR NASIONAL

COSMIC KE-3

Fakultas Kedokteran - Universitas Wijaya Kusuma Surabaya

**Tema: Tantangan Dunia Medis
dan Pendidikan Kedokteran
di Era Artificial Intelligence (AI)**

Volume 3, No 1 (2025)

[Beranda](#) / Dewan Editor

Dewan Editor

Editor in Chief

Pratika Yuhyi Hernanda, M.Sc, PhD. (Scopus Author Identifier [54892096200](#))

Editor

Dr. dr. Budhi Setiawan, M.Kes. (Scopus Author Identifier [57213297219](#))

Dr. dr. Ayling Sanjaya, Sp.A. (Scopus Author Identifier [57203983925](#))

Dr. Masfufatun, S.Si, M.Si. (Scopus Author Identifier [56038066900](#))

Dr. Dorta, Simamora, M.Si. ([Google Scholar](#))

Section Editor

Ayu Cahyani Noviani, M.KKK ([Google Scholar](#))

Noer Kumala Indahsari, S.Si., M.Si. ([Google Scholar](#))

Dr. Emilia Devi D.R., ST., MT ([Google Scholar](#))

Putu Oky Ari Tania, S.Si., M.Si ([Google Scholar](#))

Drg. Wahyuni Diah Parmasari, Sp. Ort. ([Google Scholar](#))

Steering Comite (SC)

Dr. dr. Budhi Setiawan, M.Kes. (Scopus Author Identifier [57213297219](#))

Dr. dr. Ayling Sanjaya, Sp.A. (Scopus Author Identifier [57203983925](#))

Pratika Yuhyi Hernanda, M.Sc, PhD. (Scopus Author Identifier [54892096200](#))

Beranda / Arsip / Vol 3 (2025): Edisi 2025

Vol 3 (2025): Edisi 2025



Diterbitkan: 2025-01-16

Artikel Penelitian

HUBUNGAN STATUS GIZI DENGAN DERAJAT HIPERTENSI PADA PROGRAM PROLANIS BULAN SEPTEMBER 2024 DI PUSKESMAS JATIREJO

Kiswati Rozi Permatasari, Salsabila Rahma, Bayu Prakoso, Ilham Nofa, Sukma Sahadewa
1-8



HUBUNGAN TINGKAT PENDIDIKAN DENGAN KADAR HbA1c PADA PASIEN PROLANIS DIABETES MELLITUS TIPE 2 DI PUSKESMAS DEKET LAMONGAN

Febrin Trianita Anggraeni, Roethmia Yaniari, Suhartati
9-14



PREVALENSI DAN FAKTOR-FAKTOR RISIKO INFEKSI SALURAN KEMIH PADA IBU HAMIL DI PUSKESMAS DUKUH KUPANG SURABAYA

Kresna Aditya Rukmana, Inawati, Harman Agusaputra, Budhi Setiawan
15-26



INDIKATOR PEMENDEKAN USUS MENCIT KOLITIS ULSERATIF YANG DIPULIHKAN DENGAN PEMBERIAN SCFA DAN MINYAK BIJI LABU

Gietri Indah Lestari, Putu Oky Ari Tania, Dorta Simamora, Jimmy Hadi Widjaja

27-32



PERAN KECERDASAN BUATAN DALAM KESEHATAN MASYARAKAT GLOBAL

Mohammad Yusuf Alamudi, Henry Sudiyanto, Nurwidji Nurwidji, Sulis Diana, Dhonna Anggreni, Elyana Mafticha, Eka Diah Kartiningrum, Arief Fardiansyah

33-37



EFEKTIVITAS EKSTRAK ETANOL DAUN CENGKEH (*Syzygium aromaticum*) DALAM MENGHAMBAT PEMBENTUKAN BIOFILM *Candida albicans* ATCC 14053

Firdausi Nuzula, masfufatun masfufatun

38-44



HUBUNGAN STRES AKADEMIK DENGAN GANGGUAN MAKAN (EATING DISORDER) PADA MAHASISWA FK UWKS ANGKATAN 2021 MENJELANG UJIAN SOOCA

Dani Setiawan, Dorta Simamora

55-63



PENGARUH PEMBERIAN EKSTRAK PEPEROMIA PELLUCIDA TERHADAP KADAR UREUM DAN KREATININ TIKUS WISTAR YANG MENDAPATKAN PAJANAN ASAP ROKOK

Angela Octavia, Indah Widyaningsih, Akmarawita Kadir

64-69



GAMBARAN INSIDENSI RAMPAN KARIES DENGAN KEJADIAN MALOKLUSI PADA ANAK USIA 4-8 TAHUN DI SURABAYA

Wahyuni Dyah Parmasari, Putu Agung Narendra Indria Dewi

75-80



ANALISIS MIKRO-CT SEBAGAI METODE PILIHAN TERHADAP PENELITIAN PERTUMBUHAN TULANG KRANIOFASIAL

Wahyuni Dyah Parmasari, I Gusti Wahyu Ardani, Ida Bagus Narmada, Kuntaman Kuntaman, Fourier Dzar Eljabbar Latief, Fahrishah Nurfadeliah Bahraini, Nafansya Regita Cahyani

119-123



PDF

PREVALENSI OSTEOPOROSIS PADA KEGIATAN PEMERIKSAAN KEPADATAN MASSA TULANG DI KELURAHAN PEJAGAN DAN PANGERANAN (KECAMATAN BANGKALAN)

Budhi Setiawan, Pratika Yuhyi Hernanda, Sri Lestari Utami

137-142



PDF

TINGKAT KEPUASAN PESERTA DALAM KEGIATAN PENGABDIAN MASYARAKAT “PERAN IBU TERHADAP KESEHATAN KELUARAGA BERAWAL DARI RUMAH” DI KELURAHAN KENDANGSARI SURABAYA

Agda Edita Putriska, Tasya Alfian Rahma, Ida Ayu Nandi Pusparini, Alwiyah Alwiyah, Fitrah Nirmalah, Salma Putri, Yasega Elvada, Muhammad Abi Asriansyah, Hilmi Fikri, Andikha Ardiansah, Nur Khamidah

148-153



PDF

PENGARUH EKSTRAK KAEMPFERIA PARVIFLORA TERHADAP PERTUMBUHAN BAKTERI ESCHERICHIA COLI

I Made Bimantara Febryan Putra, Lusiani Tjandra

154-159



PDF

EFEKTIVITAS TERAPI INFRAMERAH TERHADAP PENYEMBUHAN LUKA INSISI TIKUS PUTIH JANTAN

Emillia Devi Dwi Rianti, Rini Purbowati, fuad Ama

160-164



PDF

EXPLORING THE ROLE OF APOLIPOPROTEIN A AS A PREDICTIVE BIOMARKER FOR FECAL INCONTINENCE SEVERITY IN THE ELDERLY: A FOCUS ON LIPID-IMMUNE INTERACTIONS

Bryan Anna Wijaya, Peter Ian Limas, Yohanes Firmansyah, Triyana Sari, Alexander Halim Santoso

165-171



PDF

THE VISCERAL FAT SIGNIFICANTLY ASSOCIATED WITH URIC ACID LEVELS AMONG THE ELDERLY AT HANA NURSING HOME

Ayleen Nathalie Jap, Shirly Gunawan, Yohanes Firmansyah, Alexander Halim Santoso

172-178



PDF

CORRELATION BETWEEN WAIST-HIP RATIO AND WAIST CIRCUMFERENCE AS A PREDICTOR PARAMETER IN FRAMINGHAM RISK SCORE FOR HARD CORONARY HEART DISEASE AND KIDNEY FUNCTION (CARDIO-RENAL SYNDROME)

Stanislas Teguh, Andria, Ronald, Yohanes, Alexander

179-184



STRONGER HANDS, STRONGER BLADDERS: UNCOVERING THE CONNECTION BETWEEN HANDGRIP STRENGTH AND URINARY INCONTINENCE IN ELDERLY WOMEN

Fiona Valencia Setiawan, Andriana Kumala Dewi, Yohanes Firmansyah, Triyana Sari, Alexander Halim Santoso

185-190



HUBUNGAN KEJADIAN KARIES GIGI DENGAN STATUS GIZI PADA ANAK USIA 6-8 TAHUN DI SDN PUTAT JAYA I/377 SURABAYA

Luh Putu Dara Nanayanti, Theodora Theodora, Maria Juliati Kusumaningtyas, Atik Sri Wulandari

191-197



OSTEOPOROSIS PADA WARGA KELURAHAN SINGOSARI DAN DESA KARANGKIRING (KECAMATAN KEBOMAS, GRESIK) DENGAN QUANTITATIVE ULTRASOUND

Jimmy Hadi Widjaja, Sianny Suryawati, Sri Lestari Utami

198-203



Review

WAKSINASI FOODBORNE DISEASE SEBAGAI LANGKAH PENCEGAHAN RESISTENSI ANTIBIOTIK PADA ANAK

Geoffrey Christian Lo, Yohanes Firmansyah

45-54



PROBIOTIK DAN DIARE PADA ANAK

Anna Lewi Santoso, Jemima Lewi Santoso, Ayling Sanjaya

70-74



LITERATURE REVIEW: PERAN ARTIFICIAL INTELLIGENCE BERDASARKAN CHATBOT DALAM MENINGKATKAN PROMOSI KESEHATAN

Kendra Callista Satyafebrianti, Alya Shofiyah Zahira, Putri Noviana Agustina, Rafi Auliya Arbani, Muhammad Salsabeela Rusdi

81-91



HUBUNGAN KOMUNIKASI INTERPERSONAL PASIEN DAN DOKTER MELALUI KECERDASAN BUATAN (ARTIFICIAL INTELLIGENCE) : Sebuah Tinjauan Literatur

Putri Noviana Agustina, Alya Shofiyah Zahira, Kendra Callista Satyafebrianti, Rafi Auliya Arbani, Leonardo Tedjaprasadja

102-111



PERAN DOKTER DALAM MENINGKATKAN LITERASI KESEHATAN MASARAKAT MELALUI ARTIFICIAL INTELLIGENCE

Rafi Auliya, Alya Shofiyah Zahira, Kendra Callista Satyafebrianti, Putri Noviana Agustina, Fara Disa Durry

92-101



PERAN AI DALAM PEMAHAMAN DAN OPTIMALISASI KESEHATAN MENTAL PADA MASYARAKAT

Alya Shofiyah Zahira, Putri Noviana Agustina, Kendra Callista Satyafebrianti, Rafi Auliya Arbani, Hilmia Fahma

112-118



ANALISIS FAKTOR RISIKO ANEMIA PADA KEHAMILAN PENYEBAB STUNTING PADA ANAK

Karisa Betha Nanda, Susan Jocelyn Anggono, Vanessa Susanto Tjokro, Diah Arrizah Putri, Anna Lewi Santoso

124-130



A literature review PERAN MTORC1 DAN HORMON IGF-1: RISIKO OBESITAS PADA DEWASA DARI STUNTING

Susan Anggono, Vanessa Susanto Tjokro, Diah Arrizah Putri, Karisa Betha Nanda, Anna Lewi Santoso

131-136



STUDI LITERATUR: HUBUNGAN ANTARA PEROKOK PASIF DENGAN KEJADIAN KANKER PAYUDARA PADA WANITA

Umbu Reku Djawa Tana Hilungara, Titiek Sunaryati, Ira Idawati

143-147



PDF

[Focus and Scope](#)

[Editorial Board](#)

[Pendaftaran](#)

[Penyerahan Naskah](#)

[Template](#)

Tanggal Penting

Masa Pendaftaran & Penerimaan Abstrak:

Deadline 15 Oktober 2024

Batas Akhir Pengiriman Full Paper

23 November 2024

Deadline Pendaftaran Seminar

27 November 2024



Artikel Penelitian

STRONGER HANDS, STRONGER BLADDERS: UNCOVERING THE CONNECTION BETWEEN HANDGRIP STRENGTH AND URINARY INCONTINENCE IN ELDERLY WOMEN

Fiona Valencia Setiawan, Andriana Kumala Dewi*, Yohanes Firmansyah, Triyana Sari, Alexander Halim Santoso

Faculty of Medicine, University of Tarumanagara

Letjen S. Parman St No.1, RT.6/RW.16, Tomang, Grogol petamburan, West Jakarta City, Jakarta

*Email: Andrianad@stu.untar.ac.id

Abstract

Urinary incontinence (UI) is a prevalent issue among elderly women, significantly impacting their quality of life. Handgrip strength, an indicator of overall muscle strength, is understudied in its role concerning UI. Utilizing handgrip strength to measure overall muscle strength, including pelvic floor muscles, introduces a novel approach to supporting continence maintenance in elderly women. The aim is to determine the correlation between handgrip strength and UI severity among elderly women, using the International Consultation on Incontinence – Short Form (ICIQ-SF) for measurement. This study involved 74 elderly female participants at Bina Bhakti Nursing Home. Spearman's correlation was employed to analyze the relationship between handgrip strength (in kilograms), measured with the Omron Handgrip Strength device on the right hand, and UI severity was assessed through the ICIQ-SF scoring questionnaire, with scores ranging from 0 to 21. Findings indicated a statistically significant negative correlation between handgrip strength and UI severity ($r = -0.245$, $p = 0.035$). This suggests that higher handgrip strength correlates with lower UI severity. The study found a significant correlation between handgrip strength and UI in elderly women, indicating that strength-building interventions may reduce UI severity.

Keywords: elderly women, handgrip strength, ICIQ-SF, muscle strength, urinary incontinence

INTRODUCTION

As social living standards have risen, the adverse impact of incontinence on the quality of life and social dignity of older people has emerged as a significant medical and social issue. The intersection between musculoskeletal health and urinary function is an emerging area of interest in geriatric medicine, particularly in elderly women.(Li *et al.*, 2024) This association is thought to result from age-related declines in muscle mass and strength, which affect the pelvic floor muscles and contribute to urinary incontinence.(Li *et al.*, 2024; Wen *et al.*, 2023) Current assessment tools for pelvic floor muscles are often invasive, subjective, and costly. This research aims to identify a simpler, more convenient, and objective indicator to evaluate pelvic floor muscle function.(Baschung Pfister *et al.*, 2018; Li *et al.*, 2024) Chiu *et al.* found that urine stream is associated with low muscle strength not only in skeletal muscles but also in smooth muscles, such as the detrusor muscles, in a healthy male population across various ages. This finding may aid in identifying the causes of urinary dysfunction. Hand grip strength has been proposed as a biomarker for overall health.(Bohannon, 2019) This study aims to examine the correlation between handgrip strength and UI, particularly in elderly women. Most existing studies focus on other indicators of muscle strength, male participants, and unspecified age ranges.

According to Sharfina *et al.*, the highest prevalence of UI 52,17% was observed in the 50-59 age group and increased with age.(Wardani *et al.*, 2017) Examining UI by gender, women have a significantly higher prevalence compared to men.(Yavuz & Etiler, 2023) In terms of childbirth

history, the prevalence was 5% in nulliparous women, 10.25% in primiparous women, 23.61% in women with two childbirths, and 26.67% in those with three or more childbirths. The prevalence of urinary incontinence was 19.90%, with stress UI at 7.33%, urgency UI at 9.43%, and mixed UI at 3.14%. (Wardani *et al.*, 2017)

Urinary incontinence, the loss of bladder control is the unintentional release of urine due to involuntary or abnormal urine loss.(Davis *et al.*, 2020; Demaagd & Davenport, 2012) It encompasses various types of etiology stress, urge, mixed, overflow, and functional urinary incontinence. Stress urinary incontinence is a frequent condition marked by the unexpected and involuntary release of urine during actions that raise abdominal pressure, like laughing, sneezing, coughing, or physical exertion. (Lugo *et al.*, 2024) Inadequate bladder habits, such as prolonged urine retention, have detrimental effects on the strength and function of pelvic floor muscles.(Gonzalez *et al.*, 2020) Women who experience spontaneous vaginal delivery or instrumental delivery are at an increased risk of developing urinary incontinence in their later life. According to MacArthur *et al.*, urinary incontinence symptoms can persist for up to 6 years after the first birth.(MacArthur *et al.*, 2006; Saadia, 2015)

Urge incontinence often arises from issues with the detrusor muscle of the urinary bladder, a specialized smooth muscle within the bladder wall. Causes include detrusor muscle overactivity, poor detrusor compliance, and bladder hypersensitivity.(Nandy & Ranganathan, 2024) Mixed urinary incontinence involves the involuntary leakage of urine due to both stress and urge incontinence. Advanced pelvic organ prolapse (POP) leads to bladder outlet obstruction, resulting in detrusor muscle neuropathy from overstretching. This causes the bladder to overreact to neurotransmitters and lose muscle fiber synchronization, making it irritable and overactive. (Harrison *et al.*, 1990; Lo *et al.*, 2020)

Overflow incontinence occurs when the bladder is unable to empty properly, resulting in urine leakage due to weakened detrusor muscles and obstructed bladder outlets. Neurologic disorders, including spinal cord injuries, multiple sclerosis, and diabetes, can cause a neurogenic bladder. Masses in the abdomen or pelvis can cause obstruction, as can tumors from bladder cancer, which can block the urethra and lead to symptoms like dribbling after urination.(Bagnola *et al.*, 2017; Leslie *et al.*, 2014)

Functional incontinence is often observed in older adults due to environmental or physical barriers to toileting. It can result from mobility limitations, such as those following a hip fracture, which may temporarily prevent independent transfers on and off a toilet. As the underlying mobility issue improves, this condition may be resolved. In other cases, such as dementia from Alzheimer's disease or vascular disorders, patients may not recognize the sensation of a full bladder or may struggle with essential tasks, such as using the toilet.(Griebing, 2009; Leslie *et al.*, 2014)

METHOD

Research Design and Subject

To investigate the connection between handgrip and urinary incontinence, we conducted a primary analysis involving 74 elderly female participants at Bina Bhakti Nursing Home in May 2024. Participants ranged from 61-97 years old. The inclusion criteria for this study were a minimum age of 60 years and willingness to participate in the interviews. The elderly who were uncooperative, faced challenges in two-way communication, or lacked comprehension of the Indonesian language were excluded from this research.

Research Variables

The research variables included two components: handgrip and urinary incontinence. Handgrip strength, measured in kilograms, was assessed using the Omron™ Handgrip Strength dynamometer. Participants were instructed to grasp the dynamometer firmly with their left hand, hold it for a few seconds while the force was recorded, and then release it. The left hand can offer valuable insights into a person's overall strength profile, especially in cases where the dominant

hand is more frequently used in daily activities, potentially skewing results. By focusing on the non-dominant hand, we reduce this bias, providing a more balanced assessment of grip strength across individuals with different levels of handedness.(Barrios *et al.*, 2012; Incel *et al.*, 2002)

UI severity was assessed through the ICIQ-SF scoring questionnaire, with scores ranging from 0 to 21. The question items include frequency of UI, amount of leakage, the overall impact of UI, and self-diagnostic items. Higher scores indicate greater severity UI and greater impact on quality of life.(Avery *et al.*, 2004)

Statistical Analysis

Data analysis was conducted using SPSS version 26. The Kolmogorov-Smirnov test was applied to examine data distribution normality, while Spearman's Rho correlation test assessed the relationship between handgrip strength and urinary incontinence.

RESULTS AND DISCUSSION

This study examined elderly women, aged around 75 years, ranging from 61 to 97. All participants were female. On average, their handgrip strength was 8.55 kg, ranging from 1 kg to nearly 30 kg. Urinary incontinence severity was assessed using the ICIQ-SF scale, with scores ranging from 0 to 21. The average score among participants was 6.29, indicating a wide range of incontinence experiences. Such a range reflects the diversity in participants' health conditions, physical strength, and possibly lifestyle factors, all of which can impact incontinence severity.

As women get older, their urogenital system experiences several anatomical and physiological transformations. These include degeneration of muscles and axons, a decrease in bladder capacity, heightened detrusor activity, and a reduction in the strength of detrusor contractions. As a result, elderly women are more likely to experience UI. Additionally, women are more prone to this condition because of their unique anatomical, social, and cultural factors, as well as due to experiences like pregnancy, childbirth, and menopause.(Najafi *et al.*, 2022) (Table 1)

Table 1. Characteristics of Research Results

Parameter	N (%)	Mean (SD)	Med (Min-Max)
Age		74.75 (8.08)	75.50 (61 – 97)
Gender			
Female	74 (100)		
Handgrip		8.55 (5.96)	8.25 (1.00 – 29.70)
Urine Incontinence			
Yes	40 (54.1)		
No	34 (45.9)		
(ICIQ-SF)		6.29 (7.18)	3.5 (0 – 21)

With a sample size of 74, this research used the Kolmogorov-Smirnov test, which requires a minimum of 50 samples.(Mishra *et al.*, 2019) The results from the normality tests show that the data for handgrip strength and the severity of urinary incontinence do not follow a normal distribution. This conclusion is based on p-values below 0.05 from Kolmogorov-Smirnov tests. It is suggested that lower handgrip strength is associated with a higher severity of urinary incontinence among elderly women, highlighting the potential impact of muscle strength on urinary function.

Table 2. Normality Test

Parameter	Kolmogorov-Smirnov			Saphiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Urine Incontinence	0.269	74	0.000	0.810	74	0.000
Handgrip	0.113	74	0.020	0.929	74	0.000

These findings suggest that handgrip strength could serve as a convenient, non-invasive indicator of muscle health related to incontinence in elderly women. Specifically, a slight negative correlation was observed ($r = -0.245$), indicating that lower handgrip strength was associated with greater incontinence severity. In other words, individuals with weaker handgrip strength tended to experience more severe incontinence. This could make handgrip strength a simple tool for assessing health in this area, though further research is recommended to explore its potential as a measure of overall muscle health in older adults. This is different from Güzelant *et al.* which they explained found no notable difference in general muscle strength, as measured by hand grip strength, between women with POP and those without, indicating that hand grip strength might not effectively reflect pelvic floor muscle strength. Additionally, the study explored various factors that could contribute to POP, such as post-menopausal hormonal changes that reduce both muscle mass and estrogen, impacting muscle strength. Other risk factors, including obesity, were also considered and controlled for in the analysis. The accumulation of triglycerides in muscle cells can negatively affect muscle function and strength. This dysfunction may also impact the pelvic floor muscles, potentially increasing the risk of POP in obese individuals. (Aliye Yıldırım Güzelant *et al.*, 2015)

Table 3. Spearman Correlation

		Urine Incontinence	Handgrip
Urine Incontinence	Correlation Coefficient	1.000	0.245
	Sig. (2-tailed)	-	0.035
Handgrip	Correlation Coefficient	-0.245	1.000
	Sig. (2-tailed)	0.035	-

The pelvic floor muscles act like a “hammock” beneath the urethra, providing necessary support. When these muscles lose mass due to sarcopenia, urinary functions are compromised resulting in UI because the urethra cannot withstand increased bladder pressure. Sarcopenia is characterized by a progressive loss of skeletal muscle mass and strength, which can be exacerbated by factors such as aging, physical inactivity, poor nutrition, and chronic diseases. The pelvic floor muscles, including the levator ani and associated connective tissues, play a crucial role in supporting the bladder and urethra. When sarcopenia weakens and causes atrophy in these muscles, their capacity to maintain pelvic support diminishes. This reduction in muscle strength can increase bladder pressure during activities like coughing or sneezing, potentially leading to stress urinary incontinence (SUI). Another factor that sarcopenia leads to such as the production of inflammatory cytokines and oxidative stress, can cause myopathy and neurodegeneration in the bladder, ultimately leading to an overactive bladder. (Ida *et al.*, 2019; Yang *et al.*, 2021) Cytokines like tumor necrosis factor-alpha (TNF- α) and interleukin-6 (IL-6) are recognized for their role in promoting muscle wasting through several mechanisms, including the activation of the ubiquitin-proteasome system, which enhances protein degradation in skeletal muscles. These inflammatory mediators can also cause myopathy in the detrusor muscle, resulting in weakened contractions and neurodegeneration in the bladder. This neurodegeneration may disrupt the signaling pathways that regulate bladder contractions, further complicating bladder function and raising the likelihood of overactive bladder (OAB) symptoms. After accounting for various inflammation-related conditions, including heart disease, lung disease, symptomatic osteoarthritis, and diabetes, a connection was found between cytokine levels and muscle mass/strength. (Visser *et al.*, 2002) The results of this research study are similar to those of Yang *et al.*, indicating that women with sarcopenia have a

higher rate of UI. Besides that, hormonal fluctuations, particularly those occurring during menopause, can decrease muscle tone in skeletal and smooth muscles due to diminished estrogen levels. This decline adversely affects grip strength and bladder function, establishing a correlation between handgrip strength and urinary incontinence symptoms. (Maltais *et al.*, 2009; Yang *et al.*, 2021)

Exploring preventive measures like pelvic floor muscle training (PFMT) and resistance exercises can be quite advantageous for SUI, urge and mixed urinary incontinence. Research indicates that these strengthening activities not only enhance handgrip strength but also alleviate symptoms of urinary incontinence by improving the stability of the pelvic floor. Highlighting the significance of early intervention through physical therapy could play a crucial role in slowing down the progression of sarcopenia and its impact on bladder function.

CONCLUSION

This research reveals that higher handgrip strength is associated with urinary incontinence among elderly women. The results suggest that handgrip strength could serve as a simple, non-invasive measure of overall muscle health, potentially offering a practical tool to help assess and manage urinary incontinence in older women.

REFERENCES

- Aliye Yıldırım Güzelant, Nicel Taşdemir, Ayşe Banu Sarıfakıoğlu, Remzi Abalı, & Cem Çelik. (2015). *Assesment of Hand grip Strength in Patients with Pelvic Organ Prolapse*. 06(06). <https://doi.org/10.4328/JCAM.2416>
- Avery, K., Donovan, J., Peters, T. J., Shaw, C., Gotoh, M., & Abrams, P. (2004). ICIQ: A brief and robust measure for evaluating the symptoms and impact of urinary incontinence. *Neurourology and Urodynamics*, 23(4), 322–330. <https://doi.org/10.1002/nau.20041>
- Bagnola, E., Pearce, E., & Broome, B. (2017). A Review and Case Study of Urinary Incontinence. *Madridge Journal of Nursing*, 2(1), 27–31. <https://doi.org/10.18689/mjn-1000105>
- Barrios, J., Verstraeten, W., Maes, P., Aerts, J.-M., Farifteh, J., & Coppin, P. (2012). Using the Gravity Model to Estimate the Spatial Spread of Vector-Borne Diseases. *International Journal of Environmental Research and Public Health*, 9(12), 4346–4364. <https://doi.org/10.3390/ijerph9124346>
- Baschung Pfister, P., de Bruin, E. D., Sterkele, I., Maurer, B., de Bie, R. A., & Knols, R. H. (2018). Manual muscle testing and hand-held dynamometry in people with inflammatory myopathy: An intra- and interrater reliability and validity study. *PLOS ONE*, 13(3), e0194531. <https://doi.org/10.1371/journal.pone.0194531>
- Bohannon, R. W. (2019). <p>Grip Strength: An Indispensable Biomarker For Older Adults</p>. *Clinical Interventions in Aging*, Volume 14, 1681–1691. <https://doi.org/10.2147/CIA.S194543>
- Davis, N. J., Wyman, J. F., Gubitosa, S., & Pretty, L. (2020). Urinary Incontinence in Older Adults. *AJN, American Journal of Nursing*, 120(1), 57–62. <https://doi.org/10.1097/01.NAJ.0000652124.58511.24>
- Demaagd, G. A., & Davenport, T. C. (2012). Management of urinary incontinence. *P & T: A Peer-Reviewed Journal for Formulary Management*, 37(6), 345–361H.
- Gonzalez, G., Vaculik, K., Khalil, C., Zektser, Y., Arnold, C., Almario, C. V., Spiegel, B. M. R., & Anger, J. T. (2020). Women's Experience with Stress Urinary Incontinence: Insights from Social Media Analytics. *Journal of Urology*, 203(5), 962–968. <https://doi.org/10.1097/JU.0000000000000706>
- Griebing, T. L. (2009). Urinary Incontinence in the Elderly. *Clinics in Geriatric Medicine*, 25(3), 445–457. <https://doi.org/10.1016/j.cger.2009.06.004>
- HARRISON, S. C. W., FERGUSON, D. R., & DOYLE, P. T. (1990). Effect of Bladder Outflow Obstruction on the Innervation of the Rabbit Urinary Bladder. *British Journal of Urology*, 66(4), 372–379. <https://doi.org/10.1111/j.1464-410X.1990.tb14957.x>

- Ida, S., Kaneko, R., Nagata, H., Noguchi, Y., Araki, Y., Nakai, M., Ito, S., Imataka, K., & Murata, K. (2019). Association between Sarcopenia and Overactive Bladder in Elderly Diabetic Patients. *The Journal of Nutrition, Health & Aging*, 23(6), 532–537. <https://doi.org/10.1007/s12603-019-1190-1>
- Incel, N. A., Ceceli, E., Durukan, P. B., Erdem, H. R., & Yorgancioglu, Z. R. (2002). Grip strength: effect of hand dominance. *Singapore Medical Journal*, 43(5), 234–237.
- Leslie, S. W., Tran, L. N., & Puckett, Y. (2014). Urinary Incontinence. *StatPearls*. <https://www.ncbi.nlm.nih.gov/books/NBK559095/>
- Li, L., Chen, F., Li, X., Gao, Y., Wang, N., & Xu, T. (2024). Association between low handgrip strength and incontinence among Chinese middle-aged and older people: A cross-sectional study. *AGING MEDICINE*, 7(3), 360–367. <https://doi.org/10.1002/agm2.12309>
- Lo, T. S., Uy-Patrimonio, M., Kao, C., Chua, S., Huang, T.-X., & Wu, M.-P. (2020). Urodynamics mixed type urinary incontinence with advanced pelvic organ prolapse, management and outcomes. *Scientific Reports*, 10. <https://doi.org/10.1038/s41598-020-58594-3>
- Lugo, T., Leslie, S. W., Mikes, B. A., & Riggs, J. (2024). *Stress Urinary Incontinence*. StatPearls. <https://www.ncbi.nlm.nih.gov/books/NBK559095/>
- MacArthur, C., Glazener, C. M., Wilson, P. D., Lancashire, R. J., Herbison, G. P., & Grant, A. M. (2006). Persistent urinary incontinence and delivery mode history: a six-year longitudinal study. *BJOG: An International Journal of Obstetrics & Gynaecology*, 113(2), 218–224. <https://doi.org/10.1111/j.1471-0528.2005.00818.x>
- Maltais, M. L., Desroches, J., & Dionne, I. J. (2009). Changes in muscle mass and strength after menopause. *Journal of Musculoskeletal & Neuronal Interactions*, 9(4), 186–197. <http://www.ncbi.nlm.nih.gov/pubmed/19949277>
- Mishra, P., Pandey, C., Singh, U., Gupta, A., Sahu, C., & Keshri, A. (2019). Descriptive statistics and normality tests for statistical data. *Annals of Cardiac Anaesthesia*, 22(1), 67. https://doi.org/10.4103/aca.ACA_157_18
- Najafi, Z., Morowatisharifabad, M. A., Jambarsang, S., Rezaeipandari, H., & Hemayati, R. (2022). Urinary incontinence and related quality of life among elderly women in Tabas, South Khorasan, Iran. *BMC Urology*, 22(1), 214. <https://doi.org/10.1186/s12894-022-01171-9>
- Nandy, S., & Ranganathan, S. (2024). *Urge Incontinence*. StatPearls. <https://www.ncbi.nlm.nih.gov/books/NBK563172/>
- Saadia, Z. (2015). Relationship between mode of delivery and development of urinary incontinence: A possible link is demonstrated. *International Journal of Health Sciences*, 9(4), 446–452.
- Visser, M., Pahor, M., Taafe, D. R., Goodpaster, B. H., Simonsick, E. M., Newman, A. B., Nevitt, M., & Harris, T. B. (2002). Relationship of Interleukin-6 and Tumor Necrosis Factor- With Muscle Mass and Muscle Strength in Elderly Men and Women: The Health ABC Study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 57(5), M326–M332. <https://doi.org/10.1093/gerona/57.5.M326>
- Wardani, R. S., Judistiani, R. T. D., & Siddiq, A. (2017). Prevalence of Urinary Incontinence in Women aged 20–59 years in Community Dwellings. *Althea Medical Journal*, 4(2), 178–183. <https://doi.org/10.15850/amj.v4n2.1070>
- Wen, Z., Gu, J., Chen, R., Wang, Q., Ding, N., Meng, L., Wang, X., Liu, H., Sheng, Z., & Zheng, H. (2023). Handgrip Strength and Muscle Quality: Results from the National Health and Nutrition Examination Survey Database. *Journal of Clinical Medicine*, 12(9), 3184. <https://doi.org/10.3390/jcm12093184>
- Yang, S.-J., Park, J. H., Oh, Y., Kim, H., Kong, M., & Moon, J. (2021). Association of decreased grip strength with lower urinary tract symptoms in women: a cross-sectional study from Korea. *BMC Women's Health*, 21(1), 96. <https://doi.org/10.1186/s12905-021-01241-4>
- Yavuz, M., & Etiler, N. (2023). Addressing urinary incontinence by gender: a nationwide population-based study in Türkiye. *BMC Urology*, 23(1), 205. <https://doi.org/10.1186/s12894-023-01388-2>