


Experimental Dermatology

Volume 34
Number 3
March 2025




The official journal of Arbeitsgemeinschaft
Dermatologische Forschung,
European Hidradenitis Suppurativa Foundation e.V.

Editor: Akimichi Morita, Vice Director, Nagoya City
University Hospital, Professor and Chairman, Department
of Geriatric and Environmental Dermatology, Nagoya City
University Graduate School of Medical Sciences

WILEY





From lab to real-world impact

See how our Sustainable Development Goals research is shaping our future.

[Find out more](#)



Experimental Dermatology **ADF**

Editorial Information

EDITORIAL OFFICE

EXDoffice@wiley.com

EDITOR IN CHIEF

Professor Akimichi Morita

Vice Director, Nagoya City University Hospital
Professor and Chairman, Department of Geriatric and Environmental Dermatology, Nagoya City University Graduate School of Medical Sciences



Professor Morita graduated from Nagoya City University and received his M.D degree in 1989, he later received his Ph.D in basic immunology at Aichi Cancer Center. As a Humboldt Foundation fellow, he studied photobiology and photoimmunology at Duesseldorf University and underwent further training at the University of Texas Southwestern Medical Center. He was appointed Professor and Chairman of the Department of Geriatric and Environmental Dermatology at Nagoya City University Graduate School of Medical Sciences in 2003 and currently holds the position of Vice Director of Nagoya City University Hospital. He was the President of the Japanese Society of Investigative Dermatology (2018-2020) and has been involved in many other societies, such as the Japanese Society for Psoriasis Research.

ASSOCIATE EDITORS



Tom C. Chan

National Taiwan University College of Medicine (Taipei, Taiwan)

Dr. Chan is a senior consultant dermatologist at National Taiwan University Hospital. He also serves as the current Secretary-General of Taiwanese Dermatological Association. His research focuses on neuroimmune interaction and translational mechanisms of inflammatory diseases (including psoriasis, atopic dermatitis, and hidradenitis suppurativa), and cutaneous Mycobacterial infections.



Eung Ho Choi

Yonsei University Wonju College of Medicine (Wonju, Republic of Korea)

Eung Ho Choi is a Professor at the Department of Dermatology at Yonsei University Wonju College of Medicine. He obtained his MD from the Yonsei University, Wonju College of Medicine, South Korea in 1988 and PhD from the Yonsei University in 1998. He became a board-certified dermatologist in 1992. He is currently the president of the Korean Society for Investigative Dermatology (KSID). His expertise are skin barrier, atopic dermatitis, skin aging, and keratinocyte biology.



Markus Böhm

University of Münster (Münster, Germany)

Prof. Markus Böhm obtained his MD from the University of Mainz, Germany in 1991. Following his first years of residency in the Dept. of Dermatology at Free University of Berlin, Germany, he became Postdoctoral Fellow at Yale University School of Medicine, USA. After 1995 he subsequently became board-certified dermatologist, Assistant and Associate Professor at the University of Münster, Dept. of Dermatology, Germany. He is currently head of the Outpatients Department and Senior Research Scientist. His research focuses on neuroendocrinology of the skin, pigment cell biology, connective tissue and inflammatory skin diseases and hair disorders.

Choon Siew Eng

Hospital Sultanah Aminah (Johor Bahru, Malaysia)



Dr Siew Eng Choon is a senior consultant dermatologist at Hospital Sultanah Aminah, Johor Bahru, Malaysia and an Associate Professor, Clinical School Johor Bahru, Monash University, Malaysia. Dr Choon received her medical degree from the University of Malaya in Kuala Lumpur, Malaysia, and was subsequently admitted as a member of the Royal College of Physician (MRCP), London. Her main research area is chronic inflammatory skin diseases, including atopic dermatitis, psoriasis particularly pustular psoriasis, hidradenitis suppurativa and cutaneous adverse drug reactions.

**Satoshi Fukushima**

Kumamoto University (Kumamoto, Japan)

Satoshi Fukushima is a Professor at the Department of Dermatology and Plastic Surgery, Faculty of Life Sciences, Kumamoto University. His research focuses on melanoma, skin cancers, cancer immunotherapy and biomarkers.

**Nikolas Haass**

The University of Queensland (Woolloongabba, QLD, Australia)

Dr Haass is a Professor for Cutaneous Oncology and Principal Research Fellow at The University of Queensland, and a Director of International Liaison and Past President of the Australasian Society for Dermatology Research. Dr Haass obtained his MD from the University of Heidelberg, Germany in 1998, and graduated with honors with a PhD in Cell Biology in 1999. He then trained in clinical dermatology at the University of Hamburg, Germany, and moved to Philadelphia in 2003 to work as a post-doctoral fellow in Meenhard Herlyn's lab at The Wistar Institute. His research focuses on melanoma microenvironment, heterogeneity, proliferation and metastasis; 3D models and real-time 3D imaging.



Tetsuya Honda

Hamamatsu University School of Medicine (Hamamatsu, Japan)

Expertise: Cutaneous immunology, inflammatory skin diseases, lipid mediators and metabolism



Hee Young Kang

Department of Dermatology, Ajou University School of Medicine (Suwon, Republic of Korea)

Hee Young Kang is a professor and chair person of the Department of Dermatology at Ajou University School of Medicine in Suwon, South Korea. She received both her MD and PhD degrees from the Ajou University Medical School. Besides clinical dermatology, her scientific focus is in the pigmentary disorders including melasma and vitiligo and skin aging.



Dong Hun Lee

Seoul National University College of Medicine (Seoul, Republic of Korea)

Professor Dong Hun Lee is an associate professor at the Department of Dermatology, Seoul National University College of Medicine. As a physician-scientist, he received his M.D. and Ph.D. at Seoul National University. He finished his dermatology residency at Seoul National University Hospital. His translational research interests include skin aging, photodermatology, neurodermatology, epigenetic modulation, and allergic skin disorders.



Sung-Jan Lin

College of Medicine, National Taiwan University (Taipei, Taiwan)

Prof. Sung-Jan Lin is Taiwan Bio-development Foundation (TBF) Chair Professor of Department of Biomedical Engineering and Director of Research Center for Developmental Biology and Regenerative Medicine at National Taiwan University. He is also a dermatologist and Director of Center for Frontier Medicine at National Taiwan University Hospital. He is the current President of Taiwanese Society for Investigative Dermatology (TSID) and President of Taiwan Society for Stem Cell Research (TSSCR). His research interests are hair follicle biology, stem cell regulation, and regenerative biology.



Qianjin Lu

Chinese Academy of Medical Sciences and Peking Union Medical College (Peking, China)

Dr. Qianjin Lu is currently a Professor and Vice Director of Hospital for Skin Diseases (Institute of Dermatology), Chinese Academy of Medical Sciences and Peking Union Medical College; Professor and Director, Institute of Dermatology, Central South University, President of Chinese Society of Dermatology. His research focus is the epigenetics of autoimmune and inflammatory related skin diseases, including lupus, psoriasis and atopic dermatitis.



Jitlada Meephansan

Chulabhorn International College of Medicine, Thammasat University (Bangkok, Thailand)

Jitlada Meephansan is Associate Professor and Chair of Division of Dermatology at Chulabhorn International College of Medicine. Dr Meephansan obtained her MD with first class honors from Chulalongkorn University in Bangkok. She was awarded the Diploma of Dermatological Scientist by Japanese Society for Investigative of Dermatology in 2011, and has received the Young Scientist Award by Chulabhorn International College of Medicine in 2019. Her research interests are cutaneous immunology, psoriasis, hair disorders, microbiome, pruritus, and chronic inflammatory skin diseases.

Bensachee Pattamadilok

Institute of Dermatology, Department of Medical Services, Ministry of Public Health (Bangkok, Thailand)



Bensachee Pattamadilok is a senior consultant dermatologist at the Institute of Dermatology, Bangkok, Thailand. Dr. Pattamadilok graduated her Medical Degree from Mahidol University in Bangkok, Thailand. She is Board Certified in Internal Medicine, Chulalongkorn University, Board Certified in Dermatology, Mahidol University, Bangkok, Thailand, and subsequently certified as a Clinical Research Fellow, Department of Dermatology, Henry Ford Hospital and the Department of Physics, Wayne State University, Detroit, Michigan, USA. She has been selected as a Director of The Asian Society of Psoriasis. Her clinical and research interests include photodermatology, psoriasis, vitiligo, mycosis fungoides, and chronic inflammatory skin diseases. She has been the principal investigator of clinical trials investigating advanced targeted therapies in the treatment of inflammatory skin diseases.

**John Seykora**

University of Pennsylvania (Philadelphia, PA, USA)

Expertise: Keratinocyte biology, skin cancer, signal transduction, skin pathology

**Georg Stary**

Medical University of Vienna (Vienna, Austria)

Expertise: Immunodermatology, tissue-resident T cells, infectious diseases, microbiome, graft-versus-host disease, sexually-transmitted diseases

**Gang Wang**

Fourth Military Medical University, Xijing Hospital (Xi'an, China)

Dr. Gang Wang is Professor and Chair of the Dermatology Department of Xijing Hospital at the Fourth Military Medical University. He is currently the President of Chinese Dermatologist Association and Chinese Society for Investigative Dermatology. His expertise are autoimmune bullous diseases and psoriasis.

**Rei Watanabe**

Osaka University (Suita, Japan)

Dr Rei Watanabe is a professor at Osaka University, in Suita, Japan. Her research focuses on cutaneous immunology, autoimmune skin diseases, inflammatory skin diseases, skin T cells, regulatory B cells.

**Peter Wolf**

Medical University of Graz (Graz, Austria)

Dr. Peter Wolf is Professor and Chair of the Department of Dermatology and Venereology and Director of the Research Unit of Photodermatology at the Medical University of Graz, Austria. His clinical and research interests focus on photomedicine and photoimmunology. He is the principal investigator of numerous previous and ongoing clinical trials investigating phototherapeutic protocols to treat inflammatory and neoplastic skin diseases. As an active member of the PhD program of Molecular Medicine (MolMed), his translational research group addresses and elucidates the mechanisms of photo (chemo) therapy and photoimmune suppression/modulation.

**Yukie Yamaguchi**

Yokohama City University Graduate School of Medicine (Yokohama, Japan)

Dr Yamaguchi is a professor and chairman at Department of Environmental Immuno-Dermatology at Yokohama City University Graduate School of Medicine. Her research focuses on psoriasis, connective tissue diseases, cutaneous immunology, extracellular matrix.



Sook Jung Yun

Chonnam National University Medical School (Gwangju, Korea)

Sook Jung Yun is a professor and chairman of Department of Dermatology in Chonnam National University Medical School, Gwangju, Korea. She obtained her MD and PhD from Chonnam National University Medical School. Prof. Sook Jung Yun's researches focus on the melanoma, acral melanoma, skin cancers, dermatopathology, pigmentation, genetic alterations in skin cancers, and various skin manifestations in cancer patients.

Ling-juan Zhang

Xiamen University (Xiamen, China)



Dr Ling-juan Zhang is a professor in the School of Pharmaceutical Science at Xiamen University. Dr Zhang received her Ph.D. at Oregon State University and post-graduate training at the Department of Dermatology at the University of California, San Diego, where she continued as a Project Scientist and directed an NIH RO1 project. Dr. Zhang's research focuses on the innate immune mechanisms of skin host defense against bacterial infection as well as pathogenesis inflammatory skin diseases such as psoriasis. She has received several Chinese National and Fujian Province Young talent awards, and has also been elected as a board member of prestigious societies including Chinese society of Investigative Dermatology and Society of Chinese Medicine Immunology Branch.

EDITORIAL BOARD

Oleg Akilov, University of Pittsburgh (Pittsburgh)

Bogi Andersen, University of California Irvine (Irvine, CA)

Scott X. Atwood, University of California Irvine (Irvine, CA)

Zsuzsanna Bata-Csörgő, University of Szeged (Szeged)

Johann Bauer, Paracelsus Medical University (Salzburg)

Barbara Bedogni, University of Miami (Miami, FL)

Anna de Benedetto, University of Florida (Gainesville, FL)

Daniele Bergamaschi, Queen Mary University of London (London)

Mark Berneburg, University Hospital Regensburg (Regensburg)

Regina C. Betz, University of Bonn (Bonn)

Tilo Biedermann, Technical University of Munich (Munich)

Jeff Biernaskie, University of Calgary (Calgary, AB)

Daniel Bikle, University of California San Francisco (San Francisco, CA)

Ellen H. van den Bogaard, Radboud University Medical Center (Nijmegen)

Vladimir A. Botchkarev, Boston University (Boston, MA)
Joke A. Bouwstra, Leiden University (Leiden)
Nicolò Brembilla, Université de Genève (Genève)
Timo Buhl, University Medical Center Göttingen (Göttingen)
Cheng Ming Chuong, University of Southern California (Los Angeles, CA)
Antonio Costanzo, Humanitas University Milano (Rozzano-Milan)
Sandrine Dubrac, Innsbruck Medical University (Innsbruck)
Leopold Eckhart, Medical University of Vienna (Vienna)
Rüdiger Eming, Philipps-Universität Marburg (Marburg)
Sabine Eming, University of Cologne (Cologne)
Stefanie Eyerich, Technical University of Munich (Munich)
Elena Ezhkova, Icahn School of Medicine at Mount Sinai (New York, NY)
Lars French, Klinikum der Universität München (Munich)
Nicolas Gaudenzio, Université de Toulouse (Toulouse)
Richard L. Gallo, University of California San Diego (San Diego, CA)
Luis A. Garza, Johns Hopkins School of Medicine (Baltimore, MD)
Amos Gilhar, Technion- Israel Institute of Technology (Haifa)
Michel Gilliet, Lausanne University Hospital (Lausanne)
Elizabeth Grice, University of Pennsylvania (Philadelphia, PA)
Florian Gruber, AKH Vienna Dermatology (Vienna)
Frank de Gruijl, Leiden University (Leiden)
Geoffrey C. Gurtner, Stanford University (Stanford, CA)
Emma Guttman-Yassky, Icahn School of Medicine at Mount Sinai (New York, NY)
Jürgen Harder, University of Kiel (Kiel)
Matthew Hardman, University of Hull (Hull)
Cristina Has, University of Freiburg (Freiburg)
Takashi Hashimoto, Osaka City University Graduate School of Medicine (Osaka)
Tomasz Hawro, Charité Universitätsmedizin Berlin (Berlin)
Claire Higgins, Imperial College London (London)
Boris Hinz, University of Toronto (Toronto)
Ya-Chieh Hsu, Harvard University (Cambridge, MA)
Mayumi Ito, New York University (New York, NY)
Akemi Ishida-Yamamoto, Asahikawa Medical College (Asahikawa)
Ander Izeta, Donostia University Hospital Instituto Biodonostia (San Sebastián)
Francisco Jimenez, University of Las Palmas de Gran Canaria (Las Palmas de Gran Canaria)
Sarolta Kárpáti, Semmelweis Medical University (Budapest)
Kiarash Khosrotehrani, The University of Queensland (Woolloongabba, QLD)
Thomas Krieg, University of Cologne (Cologne)
Ohsang Kwon, Seoul National University College of Medicine (Seoul)
Lionel Larue, Institut Curie Centre de Recherche (Orsay)
Isabelle Caroline Le Poole, Northwestern University (Chicago, IL)
Ethan Lerner, Massachusetts General Hospital (Charlestown, MA)
Sung J. Lin, National Taiwan University (Taipei)
Fu-Tong Liu, University of California Davis (Sacramento)
Karin Loser, University of Münster (Münster)
Ralf Ludwig, University of Lübeck (Lübeck)
Thomas Magin, University of Leipzig (Leipzig)
José Carlos García-Borrón Martínez, University of Murcia (Murcia)
Kevin J. McElwee, The University of British Columbia (Vancouver, B.C.)
John McGrath, King's College London (London)
Andrew Messenger, Royal Hallamshire Hospital (Sheffield)

Michael Mildner, MUW Dermatology (Vienna, Austria)
Sarah Millar, Icahn School of Medicine at Mount Sinai (New York, NY)
Marja L. Mikkola, University of Helsinki (Helsinki)
Laurent Misery, Université Bretagne Occidentale (Brest)
Bruce A. Morgan, Harvard Medical School (Charlestown, MA)
Akimichi Morita, Nagoya City University (Nagoya)
Ulrich Mrowietz, University of Kiel (Kiel)
Motonobu Nakamura, University of Occupational and Environmental Health (Kitakyushu)
Carien Niessen, University of Cologne (Cologne)
Manabu Ohyama, Kyorin University (Tokyo)
Ryan O'Shaughnessy, Queen Mary University of London (London)
Andrey Panteleyev, Kurchatov Institute NBICS Centre Molecular Biology (Moscow)
Eva Peters, University Giessen-Marburg (Giessen)
Michael Philpott, Queen Mary University of London (London)
Mauro Picardo, San Gallicano Dermatological Institute (Rome)
Andor Pivarsci, Karolinska Institute (Stockholm)
Yuval Ramot, Hadassah-Hebrew University Medical Center (Jerusalem)
Michael Rendl, Icahn School of Medicine at Mount Sinai (New York, NY)
Michael Rosenblum, University of California San Francisco (San Francisco, CA)
Liat Samuelov, Tel Aviv Sourasky Medical Center (Tel Aviv)
Martin Schmelz, University of Heidelberg (Heidelberg)
Marlon R. Schneider, Bundesinstitut für Risikobewertung (Berlin)
Stefan W. Schneider, Universitätsklinikum Hamburg-Eppendorf (Hamburg)
George Sen, University of California San Diego (San Diego, CA)
Zhu Shen, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences (Guangzhou)
Animesh A. Sinha, University at Buffalo (Buffalo, NY)
Andrzej Slominski, University of Alabama at Birmingham (Birmingham, AL)
Eniko Sonkoly, Karolinska Institutet Department of Medicine (Stockholm)
Eli Sprecher, Tel Aviv Sourasky Medical Center (Tel Aviv)
Sonja Ständer, University of Münster (Münster)
Georg Stary, Medical University of Vienna (Vienna)
Esther von Stebut-Borschitz, University Hospital of Cologne (Cologne)
Agatha Stegemann, University of Münster (Münster)
Kerstin Steinbrink, University of Münster (Münster)
Richard A. Sturm, University of Queensland (Brisbane, QLD)
John P. Sundberg, Jackson Laboratory (Bar Harbor, ME)
Alain Taieb, Université de Bordeaux (Bordeaux)
Antonella Tosti, University of Miami (Miami, FL)
Daisuke Tsuruta, Osaka City University (Osaka)
Selma Ugurel, Universität Duisburg-Essen (Essen)
Jouni Uitto, Thomas Jefferson University (Philadelphia, PA)
Rachel Watson, University of Manchester (Manchester)
Sabine Werner, ETH Zurich (Zürich)
Sara A. Wickström, University of Helsinki (Helsinki)
Tongyu Wikramanayake, University of Miami (Miami, FL)
Miriam Wittmann, University of Leeds (Leeds)
Peter Wolf, University of Graz (Graz)
Sunny Y. Wong, University of Michigan (Ann Arbor, MI)
Gil Yosipovitch, University of Miami (Miami, FL)
Zhicao Yue, Shenzhen University College of Medicine (Shenzhen)

Zhengquan Yu, China Agricultural University (Beijing)
Patrick Zeeuwen, Radboud University Nijmegen Medical Center (Nijmegen)
Zuotao Zhao, Peking University First Hospital (Beijing)
Min Zheng, Zhejiang University (Hangzhou)
Xuejun Zhu, Peking University First Hospital (Beijing)
Paola Zigrino, University of Cologne (Cologne)
Christos Zouboulis, Dessau Medical Centre (Dessau-Rosslau)

HONORARY EDITOR

Thomas A. Luger, University of Münster (Münster)
Ralf Paus, University of Miami (Miami)



Sign up for email alerts

Enter your email to receive alerts when new articles and issues are published.

Email address

Continue

Tools



[Submit an article](#)



[Browse free sample issue](#)



[Journal Metrics](#)



[Subscribe to this journal](#)

WILEY

Discover the new
Hidradenitis Suppurativa
Knowledge Hub



Learn more

Experimental Dermatology



Experimental Dermatology: Volume 34, Issue 3

March 2025

[< Previous Issue](#) | [Next Issue >](#)

 [GO TO SECTION](#)

 [Export Citation\(s\)](#)

ISSUE INFORMATION

 [Free Access](#)

Issue Information

First Published: 06 March 2025

[PDF](#) | [Request permissions](#)

REVIEW ARTICLE

 [Open Access](#)

Skin Deep and Beyond: Unravelling B Cell Extracellular Matrix Interactions in Cutaneous Immunity and Disease

Rebecca Diehl, Stefanie Hübner, Saskia Lehr, Marta Rizzi, Kilian Eyerich, Alexander Nyström

First Published: 06 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

CONCISE COMMUNICATION

 **Open Access**

Do Melanocytes Have a Role in Controlling Epidermal Bacterial Colonisation and the Skin Microbiome?

Omera Bi, David Caballero-Lima, Stephen Sikkink, Gill Westgate, Sobia Kauser, Jacobo Elies, M. Julie Thornton

First Published: 06 March 2025


[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

RESEARCH ARTICLE

Comparing UV and Diesel Cutaneous Damage and Evaluating the Protective Role of a Topical Antioxidant Mixture Containing Vitamin C, E and Ferulic Acid

John Ivarsson, Alessandra Pecorelli, Anna Guiotto, Mariaurea Matias Souza, Hina Choudhary, Patricia Brieva, Francesca Ferrara, Giuseppe Valacchi

First Published: 10 March 2025



Effect of PM versus UV on skin oxinflammatory and structural responses. Thicker arrows indicate a stronger effect. Red colours are noxious effects while green colours indicate a protective responses by topical antioxidants (AOX) application.

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

CONCISE COMMUNICATION

Severe Darier's Disease by Mitochondrial DNA Insertion Causing Nonsense Mutations: In Silico Prediction of a Pathophysiological Mechanism to a Novel Mutation

Haruna Shintani, Yasuaki Ikuno, Hiraku Kokubu, Shino Fujimoto, Akihiko Yamaguchi, Toshifumi Takahashi, Akiko Arakawa, Yukie Kande, Hayato Naka-Kaneda, Noriki Fujimoto

First Published: 10 March 2025


[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

REVIEW ARTICLE

Human Skin Microbiota-Derived Extracellular Vesicles and Their Cosmeceutical Possibilities—A Mini Review

Thangavelu Soundara Rajan, Ramasamy Saiganesh, Madasamy Sivagnanavelmurugan, Francesca Diomede

First Published: 10 March 2025



[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

RESEARCH ARTICLE

[Open Access](#)

A Novel Ammonium Carboxylate Salt of Undecylenic Acid for the Topical Treatment of Gram-Positive and Antibiotic-Resistant Skin Infections

Alyce Mayfosh, Thomas Rau

First Published: 10 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

ABSTRACT

[Free Access](#)

51st Annual Meeting of the Arbeitsgemeinschaft Dermatologische Forschung (ADF)

First Published: 10 March 2025


CONCISE COMMUNICATION

 [Open Access](#)

Oxidative Stress and Generalised Pustular Psoriasis: Report of d-ROM Measurements in Nine Cases Including Three of Pustular Psoriasis of Pregnancy

Chisato Tawada, Yoko Ueda, Yoko Mizutani, Xiaoyu Zang, Kayoko Tanaka, Hiroaki Iwata

First Published: 10 March 2025



In pustular psoriasis, ROS-induced lipid peroxidation forms 4-HNE, triggering inflammation through NF-κB activation. This pathway, along with high IL-36 levels, recruits immune cells, creating a cycle of oxidative damage and immune response that worsens pustular psoriasis.

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

RESEARCH ARTICLE

Clinical Experience of Risankizumab in Patients With a History of Erythrodermic Psoriasis

Che-Chia Hsu, Chang-Yu Hsieh, Tsen-Fang Tsai

First Published: 12 March 2025


[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

 [Open Access](#)

Skin-Protective Performance of Alternative Stratum Corneum Formed by a Pseudo-Ceramide-Containing Steroid Lamellar Cream

Masafumi Yokota, Tomohiro Matsumoto, Akane Kawamoto, Kumiko Dojo, Sumika Toyama, Catharina Sagita Moniaga, Junko Ishikawa, Daiki Murase, Noriyasu Ota, Mitsutoshi Tominaga, Kenji Takamori

First Published: 11 March 2025



The lamellar structure-forming cream with a steroid and medical pseudo-ceramide functioned as a substitute for the stratum corneum and showed the improvement of skin barrier function and anti-inflammatory and antipruritic effects. This cream may be a promising therapeutic option for barrier-disrupted skin diseases, such as atopic dermatitis and xerosis.

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

IKZF1 and Ikaros Overexpression Results in Alopecia Areata-Like Phenotype in Mice

Yukiyasu Arakawa, Risa Tamagawa-Mineoka, Mayumi Ueta, Mari Nakanishi, Hiromi Nishigaki, Norito Katoh

First Published: 11 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

RESEARCH LETTER

 [Open Access](#)

New Characteristics of Eccrine Sweat Glands in Acquired Idiopathic Generalised Anhidrosis as Determined via Three-Dimensional Fluorescence Imaging of Cleared Skin Tissue

Satoshi Yoshida, Ryosuke Kawakami, Yosuke Niko, Yasuhiro Fujisawa, Masamoto Murakami

First Published: 12 March 2025

[Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

 **Open Access**

Multimodal Second-Harmonic-Generation, Two-Photon Excitation Fluorescence, and Brillouin Microscopy for Visualising Dermal Mechanical Properties in Ex Vivo Human Skin

Eiji Hase, Naoya Okubo, Yuki Ogura, Takeo Minamikawa, Takeshi Yasui

First Published: 12 March 2025

[Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

RESEARCH ARTICLE

Altered Sensory and Stress Responses in Atopic Dermatitis: Effects of Acute Stress on Lesional and Non-Lesional Skin

Macarena Tejos-Bravo, Dixon Cid, Fernanda Espinoza, Felipe Rojas-Thomas, Gustavo Torres, María-Laura Cossio, Arturo Borzutzky, Margarita Calvo

First Published: 17 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

Altered Arginine Metabolism Affects Proliferation and Radiosensitivity of Keloids

Wei Li, Xiaoqian Li, Yange Zhang, Baochen Zhu, Xuewen Xu, Haitao Xiao, Shuyu Zhang

First Published: 17 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

Comparative Analysis of the Skin Microbiota of Rosacea, Steroid-Induced Rosacea and Perioral Dermatitis

Akiko Mochizuki, Toshifumi Osaka, Yasuko Fukuya, Naoko Yanagisawa, Naoko Ishiguro

First Published: 17 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

Genital Psoriasis in Asians: Impact on Quality of Life and Sexual Health

Kim Yao Ong, Yee Kiat Heng, Xiahong Zhao, Hazel H. Oon

First Published: 20 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

REVIEW ARTICLE

 **Free Access**

The Impacts of Seasonal Factors on Psoriasis

Jundan Yang, Guohao Li, Lixin Yue, Erle Dang, Pei Qiao

First Published: 18 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

RESEARCH ARTICLE

 **Open Access**

The Skin Mycobiome of Patients With Atopic Dermatitis and Healthy Volunteers: A Case–Control Study

Magdalena Żychowska, Zofia Bakula, Przemysław Decewicz, Anita Hryniewicz-Gwóźdź, Mariusz Dyląg, Alina Jankowska-Konsur, Jan Gawor, Robert Gromadka, Anna Żaczek, Tomasz Jagielski

First Published: 20 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)


 **Open Access**

Effects of Emissions From Oriented Strand Board on the Development of Atopic Dermatitis Using Two Different Experimental Mouse Models

Evelyn Schneider, Katja Butter, Benjamin Schnautz, Stephanie Musiol, Johanna Grosch, Sonja

Schindela, Manuel Garcia-Käufer, Richard Gminski, Stefan Haak, Martin Ohlmeyer, Carsten B. Schmidt-Weber, Stefanie Eyerich, Julia Esser-von Bieren, Francesca Alessandrini

First Published: 20 March 2025



Higher-emitting OSB exacerbates the development of atopic dermatitis and increases lung eosinophil recruitment. Lower-emitting OSB has no or even beneficial effects in both atopic skin and lungs. Direct VOC exposures from OSB in mouse in vivo and human in vitro models

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

Hyperuricemia Exacerbates Psoriatic Inflammation by Inducing M1 Macrophage Activation and Th1 Cell Differentiation

Shu-Yi Wei, Shuang He, Xiao-Yan Wu, Yan Zhang, Ying-Ping Xu, Bin Yang, Yu-Zhe Sun

First Published: 24 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

Identification of RCC2 as a Risk Gene Associated With Basal Cell Carcinoma and Experimental Validation

Yu Zhang, Xu Han, Jiayan Ren, Man Zhu, Dake Chu, Yanmin Zhang, Qi Su, Zixi Zhang

First Published: 24 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

Semaphorin 7a Regulates the Expression of IL-4 and IL-33 in a Cell Model of Atopic Dermatitis and Is Associated With Disease Severity

Mindy Ming-Huey Guo, Kuang-Den Chen, Ho-Chang Kuo

First Published: 24 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

[Open Access](#)

How Strong Is the Link Between Merkel Cell Carcinoma and the Occurrence of Other Skin Cancer Types? A Meta-Analysis

Trairong Chokwassanasakulkit, Nigel A. J. McMillan

First Published: 26 March 2025

[Abstract](#) | [Full text](#) | [PDF](#) | [References](#) | [Request permissions](#)

CORRECTION

 **Free Access**

Correction to “The Dynamics of Circulating Tumour DNA (ctDNA) During Treatment Reflects Tumour Response in Advanced Melanoma Patients”

First Published: 26 March 2025

 **This article corrects the following: >**

The dynamics of circulating tumour DNA (ctDNA) during treatment reflects tumour response in advanced melanoma patients

Lucia Di Nardo, Laura Del Regno, Alessandro Di Stefani, Maria Mannino, Barbara Fossati, Silvia Catapano, Laura Quattrini, Yeliney Alpay, Cristina Pellegrini, Alessio Cortellini, Alessandro Parisi, Ettore Capoluongo, Chiara Autilio, Maria Concetta Fagnoli, Ketty Peris

Volume 32, Issue 10, Experimental Dermatology | pages: 1785-1793 |
First Published online: August 2, 2023

[Full text](#) | [PDF](#) | [Request permissions](#)



Sign up for email alerts

Enter your email to receive alerts when new articles and issues are published.












Email address

Continue

Tools

RESEARCH ARTICLE OPEN ACCESS

Skin-Protective Performance of Alternative Stratum Corneum Formed by a Pseudo-Ceramide-Containing Steroid Lamellar Cream

Masafumi Yokota^{1,2}  | Tomohiro Matsumoto³  | Akane Kawamoto²  | Kumiko Dojo³  | Sumika Toyama¹  | Catharina Sagita Moniaga¹  | Junko Ishikawa³  | Daiki Murase²  | Noriyasu Ota³  | Mitsutoshi Tominaga¹  | Kenji Takamori^{1,4} 

¹Juntendo Itch Research Center (JIRC), Institute for Environmental and Gender-Specific Medicine, Juntendo University Graduate School of Medicine, Chiba, Japan | ²Biological Science Research, Kao Corporation, Kanagawa, Japan | ³Biological Science Research, Kao Corporation, Tochigi, Japan | ⁴Department of Dermatology, Juntendo University Urayasu Hospital, Chiba, Japan

Correspondence: Kenji Takamori (ktakamor@juntendo.ac.jp)

Received: 28 March 2024 | **Revised:** 17 January 2025 | **Accepted:** 21 January 2025

Funding: This work was supported by Kao Corporation.

Keywords: allergen | ceramide | proteases | pruritus | skin barrier function

ABSTRACT

Ceramides in the stratum corneum (SC) are important for epidermal barrier function. We previously developed a synthetic pseudo-ceramide for medical (SPCM)-containing steroid cream [SPCM (+)]. This cream forms films on the skin surface and exerts anti-inflammatory effects through steroids. However, the preventive effects of this cream on the disruption of the skin barrier remained unclear. Therefore, in this study, we aimed to evaluate the protective role of SPCM (+) cream against atopic dermatitis (AD)-associated protease allergens on the skin in recovery from barrier-broken skin. We used three-dimensional (3D) skin and mouse models of disrupted skin barriers to evaluate the protective effect of SPCM (+) cream against V8 protease produced by *Staphylococcus aureus*. In NC/Nga mice with itching caused by living mites, SPCM (+) cream was repeatedly applied once a day for 2 weeks, and scratching behaviour was assessed every week using the MicroAct system. In the 3D skin model, the SPCM (+) cream directly blocked SC degradation by V8 protease of *S. aureus* and suppressed the expression of interleukin-36 gamma. The application of SPCM (+) cream to mite-parasitised mice suppressed scratching, reduced elevated activity of skin proteases, and inhibited upregulation of thymic stromal lymphopoietin. These beneficial effects of SPCM (+) cream were not observed with steroid creams without SPCM. These results suggest that the SPCM (+) cream is effective in relieving inflammation and itching by protecting the skin from proteases and allergens through its lamellar structure. This cream may be a promising treatment option for skin barrier disorders including AD and xerosis.

1 | Introduction

The epidermis protects the body from environmental stress triggered by physical, chemical, and pathological factors and acts as a barrier to prevent water loss from the body. Intercellular lipids, known as ceramides, play an essential role in the skin barrier function. Ceramides contribute to the integrity of the epidermal barrier by organising lipid multilayers (lamellar) together with

cholesterol and free fatty acids [1]. Shortage and aberrant profiles of ceramides in the stratum corneum (SC) are closely linked with the pathogenesis of atopic dermatitis (AD) and psoriasis, which are mainly characterised by impairment of the skin barrier [2, 3].

Improving skin barrier function is considered very important in AD because penetration by external irritants causes the

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Experimental Dermatology* published by John Wiley & Sons Ltd.

onset or relapse of inflammation [4]. Harmful components commonly associated with AD exacerbation are abundant in the environment, and allergens derived from *Staphylococcus aureus*, house dust mites, and pollen are well known [5]. Most allergens possess proteolytic activity (protease allergens) and can disrupt the epidermal barrier. Moreover, protease allergens and mechanical barrier damage, such as that caused by scratching, collaboratively facilitate epicutaneous sensitization, skin inflammation, such as AD, and atopic march including asthma [6]. Consequently, ameliorating impaired skin barrier function and protecting skin from allergens are important for skin and systemic health.

The topical application of ceramides to skin lacking intercellular lipids and with an impaired skin barrier restores barrier function and the water content of the SC [7, 8]. However, extraction from living organisms or the chemical synthesis of natural ceramides for skin application poses serious safety and cost related challenges [9]. Natural ceramides derived from biogenic sources are primarily extracted from bovine pituitary (bovine pituitary extracts, BPE), and prion-contaminated BPE can transmit bovine spongiform encephalopathy (BSE) [10]. Although synthetic ceramides do not carry the risk of infectious diseases like BSE, producing skin-equivalent ceramides is highly expensive (2000–several hundred thousand €/kg) [11]. The commercial use of natural ceramides derived from plants and yeast has been growing in recent years; however, issues regarding production, such as low yields and difficulties in extraction and purification, need to be resolved [12, 13].

Therefore, pseudo-ceramides mimicking skin-identical ceramides, such as PC-104 {N,N'-(2-hydroxy-1,3-propanediyl)bis[N-(2-hydroxyethyl)hexadecanamide]} and Bio391 [N-(2-hydroxyethyl)-2-pentadecanoyl hexadecanamide], have been developed and tested. These pseudo-ceramides are relatively cheap to produce and less-toxic; in addition, creams and lotions containing pseudo-ceramides effectively improve skin barrier function in cases of xerosis and AD, as well as those containing skin-derived ceramides [9, 14, 15]. SPCM [N-(3-hexadecyloxy-2-hydroxypropyl)-N-2-hydroxyethyl hexadecanamide] is one of the synthetic pseudo-ceramides, and mimics naturally occurring ceramide type 2 (NS) [16]. Supplementation with SPCM restores barrier integrity and water-holding properties of the SC, thereby exhibiting therapeutic efficacy against experimental barrier-disrupted and dry skins of patients with AD [16, 17]. Recently, we have developed an oil-in-water cream by combining a steroid (prednisolone valerate acetate, PVA) with SPCM. This SPCM-containing steroid cream [SPCM (+) cream] has improved disease severity to the same extent as that of commercial steroid creams after 2 weeks of application in patients with AD. Notably, SPCM (+) cream substantially restores skin properties, such as moisture content of the SC and transepidermal water loss (TEWL), more quickly than the steroid cream lacking SPCM [18]. Formulations with multilamellar structures behave like SC with appropriate moisture permeability while limiting substance penetration [19, 20]. These findings imply that the protective effect of the lamellar structure may play an important role in the efficacy of SPCM (+) cream and in replenishing SPCM and anti-inflammatory effects of steroids. However, the mechanism by which the alternative SC formed by SPCM (+) cream contributes to skin restoration is not well understood.

In the present study, we focused on the harmful effects of AD-associated protease allergens on skin and evaluated the potential of SPCM (+) cream to recover barrier-broken skin, similar to that of AD, using in vitro and in vivo models. Herein, we describe how the SPCM (+) cream suppresses the expression of cytokines, such as interleukin-36 gamma (IL-36γ) and thymic stromal lymphopoietin (TSLP), by protecting skin from proteases, as an alternative SC, and alleviates skin inflammation and itching.

2 | Methods

2.1 | Test Creams

In this study, synthetic pseudo-ceramide of medical-use grade was used. A synthetic pseudo-ceramide for medical (SPCM) was provided by Kao Corporation (Tokyo, Japan). The chemical structure of SPCM is shown in Figure S1. The SPCM (+) cream was an oil-in-water formulation containing 0.15% prednisolone valerate acetate (PVA) and 3% SPCM. The oil-in-water formulation was prepared by mixing the aqueous and oil phases at 80°C–85°C and cooling to room temperature. Other components included cetyl alcohol, stearyl alcohol, polyoxyethylene sorbitan monostearate, sorbitan monostearate, glycerin, and water. The control cream [SPCM (–) cream] was formulated by replacing SPCM with water. The cream containing no SPCM or PVA was designated as the vehicle cream (Vehicle).

2.2 | Reagents

Mouse anti-CDSN antibody was purchased from Abcam (Cambridge, UK). Alexa Fluor 488-conjugated donkey anti-mouse IgG antibody, Dulbecco's Modified Eagle Medium (DMEM) high glucose, and Ham's F-12 Nutrient Mix were purchased from Thermo Fisher Scientific (Waltham, MA, USA). *S. aureus* V8 protease was purchased from FUJIFILM Wako Pure Chemical (Osaka, Japan) and dissolved in Dulbecco's phosphate-buffered saline (PBS, Thermo Fisher Scientific) for use.

2.3 | Structural Observation by Transmission Electron Microscopy

The test cream was applied onto a polyethylene terephthalate film using a coater (thickness, 120 μm). After 3 h of drying, they were cut into 5-mm squares, stained, and fixed with osmium tetroxide for 24 h at 23°C. A Leica UC-6 cryo-ultramicrotome (Leica Microsystems, Wetzlar, Germany) was used to obtain ultrathin sections of 50–100 nm thickness at –80°C. The sections were collected in droplets of frozen 2.3 M sucrose, placed on a grid (200 mesh with film without carbon reinforcement), washed and dried. Transmission electron microscopy was performed using an H-7650 (Hitachi, Tokyo, Japan) at an accelerating voltage of 80–100 kV. A charge-coupled device (1024 × 1024 pixels) was used for detection.

2.4 | Cell Culture

A three-dimensional (3D) human epidermal model (LabCyte EPI-MODEL12 6D) was obtained from Japan Tissue Engineering

Co. Ltd. (Aichi, Japan). Cells were pre-incubated overnight with culture media provided by the manufacturer at 37°C in a humidified atmosphere of 95% air and 5% CO₂. After preincubation, the medium was replaced with conditioned medium consisting of a 3:1 mixture of DMEM and Ham's F-12 medium. Each cream was applied topically to the 3D skin model and dried for 30 min. Subsequently, filter paper absorbed with 60 µL PBS or 1% (w/v) V8 protease solution was overlaid on the coated cream, and cells were harvested after 3 h.

2.5 | Immunostaining

Frozen blocks were prepared by embedding unfixed skin tissues in Super Cryoembedding Medium (Leica Microsystems, Wetzlar, Germany). Next, 6-µm-thick cryosections were cut using a Cryostat Microm HM550 (Thermo Fisher Scientific), dried, and fixed in 4% paraformaldehyde for 4 h. The sections were blocked with Protein Block Serum-Free Ready-to-use (Agilent Technologies, Santa Clara, CA, USA) for 10 min at 25°C and then incubated with primary antibody at 4°C overnight. The secondary antibody and 4',6-diamidino-2-phenylindole (DAPI) solution (Thermo Fisher Scientific) were added to the sections and incubated at 25°C for 30 min, followed by the addition of Fluoromount-G (Southern Biotech, Birmingham, AL, USA). Antibodies and DAPI were diluted using Can Get Signal Solution A (TOYOBO, Osaka, Japan). Immunofluorescent images were captured using a confocal fluorescence microscope LSM710 and imaging software ZEN v.3.1 (blue edition) (Carl Zeiss Microscopy, Jena, Germany).

2.6 | mRNA Extraction and Quantitative Real-Time Polymerase Chain Reaction (PCR)

Total RNA was extracted from cultured cells and mouse tissues using an RNeasy Mini Kit (QIAGEN, Venlo, the Netherlands) and reverse-transcribed using a High-Capacity RNA-to-cDNA kit (Thermo Fisher Scientific). Quantitative real-time PCR was performed on a 7500 Fast Real-Time PCR System (Thermo Fisher Scientific) using TaqMan Fast Universal PCR Master Mix (Thermo Fisher Scientific). The following human and mouse genes were analysed using TaqMan Gene Expression Assays (Thermo Fisher Scientific): human *RPLP0* (Hs00420895_gH), mouse *Rplp0* (Mm00725448_s1), human *IL36G* (Hs00219742_m1), human *CXCL8* (Hs00174103_m1), human *MMP9* (Hs00957562_m1), human *DEFB4B* (Hs00823638_m1), and mouse *Il36g* (Mm00463327_m1).

2.7 | RNA Sequencing (RNA-Seq)

Sequencing libraries were prepared using an Ion AmpliSeq Transcriptome Human Gene Expression Kit (Thermo Fisher Scientific). Briefly, RNA was reverse-transcribed using a SuperScript VILO cDNA Synthesis Kit (Thermo Fisher Scientific). cDNA was ligated with Ion Xpress Barcode adaptors and purified using AMPure XP beads (Beckman Coulter, Brea, CA, USA). A quality check of the library was performed using a High Sensitivity D1000 ScreenTape on an Agilent 4200 TapeStation. Each library was quantified using an Ion Library

TaqMan Quantitation Kit (Thermo Fisher Scientific) and diluted to 100 pM prior to template preparation. Templates were prepared and loaded onto an Ion 540 chip in an Ion Chef System (Thermo Fisher Scientific). RNA-seq was performed using an Ion S5 XL System (Thermo Fisher Scientific).

2.8 | RNA-Seq Data Analysis

Read count data were generated using the AmpliSeq RNA plug-in of the Ion Torrent Suite Software v.5.14.0 (Thermo Fisher Scientific). Differentially expressed genes (DEGs) were analysed using integrated Differential Expression and Pathway analysis (iDEP) v.0.94. iDEP is a web-based tool for analysing RNA-seq data based on the R/Bioconductor packages [21]. DEGs were defined with a false discovery rate < 0.05, and fold-change > 1.5.

2.9 | Enzyme-Linked Immunosorbent Assay

Whole mouse skin was homogenised and sonicated using a CelLytic MT Cell Lysis Reagent (Sigma-Aldrich, St. Louis, MO, USA) supplemented with a protease/phosphatase inhibitor cocktail (Cell Signaling Technology, Danvers, MA, USA). After centrifugation (15 000 × g for 15 min at 4°C), the supernatants were collected. Concentrations of IL-36γ and TSLP were determined according to the manufacturer's instructions. Protein quantity was determined using a BCA protein assay kit (Thermo Fisher Scientific). The amounts of cytokines were adjusted according to protein quantity of the supernatants.

2.10 | Mice

Male ICR mice (eight-week-old) and male NC/Nga mice (eight-nine-week-old) were purchased from Japan SLC (Shizuoka, Japan). These mice were kept under controlled conditions with a 12-h light/12-h dark cycle and a steady temperature of 23°C ± 2°C and had ad libitum access to water and food. All animal experiments were approved by the Animal Ethics Committee of Kao Tochigi Institute (approval number: F17079-0001, K20037-01), Juntendo University (approval number: 2022112), and Japan SLC (approval number: E77-8111).

2.11 | Skin Barrier Disruption and Protease-Challenge Model

Three days after shaving the rostral backs of mice, cotton soaked in a mixture of acetone and diethyl ether (1:1) was applied to the shaved backs for 15 s, immediately followed by soaking with distilled water for 30 s [acetone-ether-water (AEW) treatment]. The treatment was repeated twice daily at intervals of more than 6 h for 2 days. AEW treatment was followed by cream application and exposure to V8 protease. 40 µL of each cream were applied to the shaved area (2 × 2 cm) and dried for 1 h. Using a syringe equipped with a nasal nebulizer (Fuji-Medical Corporation, Tokyo, Japan), 0.02% (w/v) V8 protease solution was sprayed onto the cream-treated area and dried for at least 30 min. These processes were continued for 3 days. All treatments

were performed under isoflurane anaesthesia (Viatris Inc., Canonsburg, PA, USA).

2.12 | Measurement of TEWL

TEWL was evaluated using a Tewameter TM300 (Courage and Khazawa, Cologne, Germany). Measurements were conducted by placing the device on the surface of the back skin under isoflurane anaesthesia.

2.13 | Mite-Infested AD Model

NC/Nga mice develop AD-like dermatitis under conventional circumstances, but not under specific pathogen-free (SPF) conditions [22]. In this study, dermatitis was induced in mice parasitised with *Myobia musculi* in a conventional environment to increase the efficiency of this model [23] and to evaluate the protective effect of each cream against mite proteases. Each cream (50 mg) was topically applied to the shaved back (2 × 2 cm) and ears once daily for 2 weeks. Mice were anaesthetised with isoflurane (Viatris Inc.) during the experimental procedure.

2.14 | Quantification of Scratching Behaviours

Scratching behaviour was automatically and objectively recorded using MicroAct (Neuroscience, Tokyo, Japan) as previously described [24]. Briefly, a Teflon-coated magnet (1.0 mm in diameter, 3.0 mm in length) was subcutaneously implanted into the hind paws of isoflurane-anaesthetised mice 3 days before measurement. Each mouse was placed in an observation chamber surrounded by a circular coil containing food and water. Scratching behaviour was defined as electric current in the coils induced by the movement of magnets in the hind paws. The analysis parameters for detecting scratch motions are listed in Table S1. The number of scratching events was counted over 22 h.

2.15 | Skin Protease Activity Assay

Protease activity in the skin was assessed using an EnzCheck Protease Assay Kit (Thermo Fisher Scientific) consisting of casein derivatised with a fluorescent dye (casein-FL). Although undigested casein-FL is quenched, casein-FL cleaved by proteases can fluoresce. The lesion skin of NC/Nga mice was dissected, homogenised, and sonicated in a buffer supplied by the manufacturer. The supernatant was separated by centrifugation (15000 × g for 15 min at 4°C) and used for the assay. Protease activity was measured immediately after mixing the supernatant with casein-FL over a period of 60 min. Fluorescence was measured at 485 nm (excitation) and 530 nm (emission) using a plate reader SH-9000 (Hitachi High-Tech Corporation, Tokyo, Japan).

2.16 | Statistical Analysis

Data are presented as the mean ± standard error (SEM). Student's *t*-test and one-way analysis of variance with Dunnett's

or Tukey's multiple comparison tests were used for statistical analyses. All analyses were performed using GraphPad Prism v.8 (GraphPad Software, La Jolla, CA, USA).

3 | Results

3.1 | Lamellar Structure Formed by the SPCM (+) Cream Mitigated Damage to the Epidermal Barrier by V8 Protease

Lamellar structure formed by the SPCM (+) cream was confirmed using transmission electron microscopy (Figure 1A). However, no lamellar structure was observed for the SPCM (−) cream (data not shown). We conducted experiments using a 3D skin model (Figure 1B) to evaluate the protective effect of SPCM (+) cream against V8 protease produced by *S. aureus* [25]. V8 protease is involved in the pathogenesis of AD by causing disruption of the skin barrier and inflammation [26, 27]. The influence of V8 protease on the SC was evaluated by immunostaining based on the fluorescence intensity of corneodesmosin (CDSN) (Figure 1C). Exposure of the 3D skin model to V8 protease without any treatment resulted in the disappearance of the SC (Figure 1D). In the SPCM (−) cream-treated group, the CDSN signal was attenuated, and the surface of the SC was partially stripped, indicating that the SC structure was considerably damaged (Figure 1E). No damage to the SC was observed in the SPCM (+) cream-treated group, which was comparable to that observed in the V8 protease-untreated group (Figure 1F).

3.2 | IL-36γ, an Inflammatory Cytokine, Reflected Skin Response to V8 Protease

Next, we analysed gene expression in 3D skin models using RNA-seq. DEGs analyses between PBS- and V8 protease-treated samples revealed that *IL36G* was markedly upregulated by V8 protease (Figure 2A). Analysis of DEGs between groups with and without SPCM (+) cream treatment under V8 protease stimulation showed that *IL36G* expression was markedly decreased in the SPCM (+) cream-treated group (Figure 2B). We further validated the RNA-seq data using real-time quantitative PCR and obtained data consistent with that of the RNA-seq analysis (Figure 2C). In addition to *IL36G*, C-X-C motif chemokine ligand 8 (*CXCL8*), matrix metalloproteinase 9 (*MMP9*), and defensin beta 4B (*DEFB4B*) were identified as candidate markers for skin response to V8 protease (Figure S2). However, these are downstream genes of the IL-36γ signalling pathway [28, 29] and *CXCL8* is deleted in the mouse and rat genomes [30], limiting further investigation. For these reasons, we selected the IL-36γ expression as a suitable marker for evaluating skin reactions to V8 protease.

3.3 | The SPCM (+) Cream Conferred Barrier Function on the Impaired Skin

Both AEW treatment and the application of V8 protease break the epidermal barrier in mice [31, 32]. Therefore, we designed




FIGURE 1 | Lamellar structure of SPCM-containing cream and its protective effect against V8 protease. (A) Representative transmission electron microscopic images of lamellar structure formed by the SPCM (+) cream. Scale bar, 100 nm. (B) Scheme of the experiment using a reconstructed human epidermis model. (C–F) Representative fluorescence images of 3D skin models stained with anti-CDSN antibody (SC marker, green) and DAPI (nuclear marker, blue). Scale bars, 100 μ m. CDSN, corneodesmosin; NT, untreated samples; PVA, prednisolone valerate acetate; SC, stratum corneum; SPCM (-), PVA cream without SPCM; SPCM (+), SPCM-containing PVA cream; SPCM, synthetic pseudo-ceramide for medical.




FIGURE 2 | IL-36 γ is a marker of V8 protease-induced SC. (A, B) Volcano plot of DEGs in the V8 protease-treated group vs. control group (A) and V8 protease + SPCM (+)-treated group vs. V8 protease-treated group (B). Upregulated genes are towards the right, and downregulated genes are towards the left [$n = 3$, Benjamini–Hochberg false discovery rate (FDR) < 0.05, fold change (FC) > 1.5]. (C) IL36g mRNA expression was validated by quantitative real-time PCR. $n = 3$. Data represent mean \pm SEM. **** $p < 0.0001$, ** $p < 0.01$, Dunnett's test (vs. NT under V8 protease challenge). IL-36 γ , interleukin-36 γ ; NT, untreated samples; SC, stratum corneum; SPCM (+), SPCM-containing PVA cream; SPCM, synthetic pseudo-ceramide for medical.

an experimental model that combined AEW treatment and V8 protease exposure to investigate the efficacy of SPCM (+) cream on barrier-impaired skin (Figure 3A). TEWL was significantly reduced in the SPCM (+) cream-treated group compared to that in groups after AEW treatment (Day 2) and V8 protease application (day 4) ($p = 0.0483$) (Figure 3B). IL-36 γ levels in the V8 protease-sprayed and unsprayed areas reflected the effect of

V8 protease on murine skin, which was similar to that of the 3D skin model (Figure S3). To determine whether protection against V8 protease was associated with the recovery of TEWL, we compared the IL-36 γ expression among groups. IL-36 γ expression was suppressed in the SPCM (+) cream-treated group compared to that in the untreated and SPCM (-) cream-treated groups (Figure 3C,D).




FIGURE 3 | The SPCM-containing cream serves as a substitute of SC on the disrupted skin. (A) Scheme of the experiment of skin barrier disruption and protease challenge model. (B) Changes in skin barrier function (TEWL) of the SC before and after application of the test cream. $n = 8$. Data represent mean \pm SEM. * $p < 0.05$, Paired t -test. (C, D) The mRNA (C) and protein (D) expression of IL-36 γ after V8 protease stimulation along with treatment of the test cream. $n = 8$. Data represent mean \pm SEM. * $p < 0.05$, Tukey's test. IL-36 γ , interleukin-36 γ ; n.s., not significant; NT, untreated samples; SC, stratum corneum; SPCM (-), PVA cream without SPCM; SPCM (+), SPCM-containing PVA cream; SPCM, synthetic pseudo-ceramide for medical; TEWL, transepidermal water loss.

3.4 | Regular Use of the SPCM (+) Cream Suppressed Scratching Behaviour in AD Mice

Similar to V8 protease, mite proteases adversely affect skin [5]. As allergens, they also contribute to the development of AD, and their relationship with pruritus has been reported [33]. Therefore, we hypothesised that the SPCM (+) cream would prevent itching triggered by mite-derived substances. Using NC/Nga mice with itching caused by living mites, scratching behaviour was assessed every week, and each cream was repeatedly applied once a day for 2 weeks (Figure 4A). The frequency of scratching bouts did not change in the Vehicle-treated group, whereas scratching behaviours significantly decreased in the SPCM (+) cream-treated group from the first week onwards. Treatment with the SPCM (-) cream also reduced scratching behaviour, albeit to a lesser extent than that by the SPCM (+) cream-treated group (Figure 4B). Comparison among the groups at each time point showed that the application of SPCM (+) cream only suppressed itching in the first week, and this trend was more evident in the second week (Figure 4C).

3.5 | Protease Activity and TSLP Expression in the Skin of AD Mice Were Reduced by Treatment With the SPCM (+) Cream

Mite-derived proteases evoke itching by activating protease-activated receptor 2 (PAR2) [34]. Therefore, we focused on skin protease activity to determine the mechanism underlying the inhibition of itching by the SPCM (+) cream. Skin protease activity

increased in the Vehicle-treated mice relative to that in the SPF mice. The SPCM (+) cream-treated group displayed a significant reduction in skin protease activity (Figure 4D). TSLP is a representative itch mediator secreted by keratinocytes, and its release is attributed to PAR2 activation in keratinocytes [35]. The SPCM (+) cream-treated group exhibited the lowest TSLP level among all groups (Figure 4E).

4 | Discussion

In the present study, using 3D skin models, dry skin and AD model mice, we found that the topical application of the SPCM (+) cream to the skin with weakened barrier function suppressed inflammation (increased levels of IL-36 γ and TSLP), disruption of skin barrier function (increase in TEWL), and itching (increased number of scratching bouts), by preventing the entry of foreign proteases and living mites into the body. Because we observed that the SPCM (-) cream treatment partially suppressed the degradation of SC by V8 protease, the SPCM (-) cream can be expected to have a certain protective effect. However, the CDSN signal in the SPCM (-) cream-treated group was obviously attenuated and a part of the SC detached compared to observations in the SPCM (+) cream-treated group. These findings indicate that mixing the cream with SPCM provided protective effects other than the ability to cover the skin surface. The dried film of SPCM (+) cream formed regular layers at approximately 10-nm intervals. Periodic lamellar organisation of intercellular lipids of the human SC with an interval of 6–13 nm were very similar with the structure formed by the SPCM (+) cream [36]. The lamellar phase present in intercellular lipids

the SPCM (+) cream are expected to be effective against *S. aureus* proteases, the SPCM (+) cream may be useful for patients with AD under skin conditions where *S. aureus* is likely to colonise the deep cutaneous layers, to maintain skin health.

In the skin barrier disruption and protease challenge models, the SPCM (+) cream significantly reduced TEWL, whereas the SPCM (–) cream did not affect TEWL. A moist environment facilitates epidermal turnover, thereby quickly normalising the function of SC. The film formed by the SPCM (+) cream exhibits high moisture retention and occlusive properties, and these effects are at least partly attributed to its lamellar structure on the skin [20]. AEW treatment transiently elicits skin dryness, and the water-retention capability of the SPCM (+) cream is expected to restore the epidermal barrier. Compared to that of the SPCM (–) cream-treated group, IL-36 γ expression in the skin was downregulated in the SPCM (+) cream-treated group. Aberrantly enhanced IL-36 γ signalling exacerbates tissue damage, and downregulation of IL-36 γ function promotes wound repair [40, 41]. Therefore, suppression of IL-36 γ expression by the SPCM (+) cream may partly contribute to improving the skin barrier. An SPCM lamellar cream designed with the same strategy was reported to positively change the ceramide profile of the SC and improve TEWL [42]. According to this study, the penetration of SPCM from the SPCM (+) cream into the damaged SC may also have contributed in part to the skin barrier restoration. However, the primary mechanism of skin barrier recovery is unknown, thus necessitating further detailed research, such as measurement of SPCM penetration into the SC.

Behavioural analyses showed that treatment with the SPCM (+) cream significantly inhibited scratching behaviour in mite-infested AD model mice. Skin disorders with chronic itching, such as AD, psoriasis, and xerosis, lead to a vicious itch-scratch cycle, in which frequent scratching aggravates skin symptoms and further intensifies itching [43]. To overcome this negative cycle, one rational therapeutic strategy to reinforce skin integrity is to reduce scratching. Although PVA included in the SPCM (+) and SPCM (–) creams is a relatively weak steroid, its usefulness in relieving pruritus in AD is well established [44]. From permeability tests using porcine skin, we verified that transdermal permeability of PVA was comparable between the SPCM (+) and SPCM (–) creams at 24h after treatment. Therefore, the antipruritic effect of SPCM (+) cream was not dependent on the anti-inflammatory effect of steroids but was largely attributed to characteristic features of the physical properties of SPCM, including its lamellar structure. Epidermal barrier dysfunction is observed in the skin of patients with AD in lesion and non-lesion areas [45]. Therefore, proactive therapy using moisturisers and low-dose topical anti-inflammatory drugs is important for maintaining a healthy state of the skin barrier in the early stages before skin lesions become apparent. Therefore, PVA-containing creams that provide alternative SC by SPCM may be promising tools for proactive therapy.

Protease activity in the skin of AD mice was significantly reduced in the SPCM (+) cream-treated group. PAR2 is activated by proteolytic cleavage of its extracellular N-terminus and is associated with eczema-like inflammation and pruritus [46]. Epidermal protease activity is derived from endogenous or exogenous sources. Kallikrein 5 (KLK5) is a major endogenous protease that is involved in SC formation. KLK5 is upregulated

in the SC of patients with AD, and its overexpression results in increased scratching via PAR2 activation [47, 48]. However, in the present study, mRNA expression of KLK5 in skin was not altered by the application of SPCM (+) cream (Yokota et al. unpublished observations). The main allergens from house dust mites possess serine or cysteine protease activity [5]. These exogenous proteases penetrate the skin by degrading the epidermal barrier and inducing itching via PAR2 stimulation [34]. These results imply that the SPCM (+) cream protects the epidermis from protease allergen invasion, thereby inhibiting skin protease activity. This idea was further supported by the fact that the level of TSLP that is transcribed downstream of PAR2 signalling was relatively low in the SPCM (+) cream-treated group.

In conclusion, the lamellar structure-forming SPCM (+) cream may act as a substitute for the SC as it enhances skin barrier function and anti-inflammatory and antipruritic effects. The SPCM (+) cream has promising therapeutic potential to prevent barrier-disrupted skin diseases such as AD and xerosis.

Author Contributions

Conceptualization: Masafumi Yokota, Tomohiro Matsumoto, Akane Kawamoto, Kumiko Dojo. Methodology: Masafumi Yokota, Tomohiro Matsumoto, Akane Kawamoto, Kumiko Dojo. Validation: Masafumi Yokota, Tomohiro Matsumoto, Akane Kawamoto, Kumiko Dojo, Sumika Toyama, Catharina Sagita Moniaga. Formal analysis: Masafumi Yokota, Tomohiro Matsumoto, Akane Kawamoto. Investigation: Masafumi Yokota, Tomohiro Matsumoto, Akane Kawamoto, Kumiko Dojo, Sumika Toyama, Catharina Sagita Moniaga. Resources: Noriyasu Ota, Mitsutoshi Tominaga, Kenji Takamori. Data curation: Masafumi Yokota. Writing – original draft: Masafumi Yokota; Writing – review and editing: Masafumi Yokota, Kenji Takamori. Visualisation: Masafumi Yokota, Tomohiro Matsumoto, Akane Kawamoto. Supervision: Junko Ishikawa, Daiki Murase. Project administration: Noriyasu Ota, Mitsutoshi Tominaga, Kenji Takamori. Funding acquisition: Noriyasu Ota. All authors read and approved the final manuscript.

Acknowledgements

We thank Tsuyoshi Seki, Azumi Nagasawa, Shotaro Ito, Michiyo Sasajima, and Haruka Fujikawa at Health and Wellness Products Research, Kao Corporation for their technical support and helpful discussions. We would also like to thank Japan SLC for their assistance with the animal experiments. The authors are grateful to Editage (www.editage.jp) for English language editing.

Conflicts of Interest

MT and KT received joint research funds from Kao Corporation. The other authors declare no conflicts of interest.

Data Availability Statement

RNA-seq data have been deposited in the DNA Data Bank of Japan (DDBJ) Sequenced Read Archive (<https://www.ddbj.nig.ac.jp/>) as follows: BioProject: PRJDB16927, Run: DRR512897-DRR512905. Other data related to this article are available from the corresponding author on reasonable request.

References

1. Y. Uchida and K. Park, “Ceramides in Skin Health and Disease: An Update,” *American Journal of Clinical Dermatology* 22, no. 6 (2021): 853–866.

2. J. Ishikawa, H. Narita, N. Kondo, et al., "Changes in the Ceramide Profile of Atopic Dermatitis Patients," *Journal of Investigative Dermatology* 130, no. 10 (2010): 2511–2514.
3. U. Yokose, J. Ishikawa, Y. Morokuma, et al., "The Ceramide [NP]/[NS] Ratio in the Stratum Corneum Is a Potential Marker for Skin Properties and Epidermal Differentiation," *BMC Dermatology* 20, no. 1 (2020): 6.
4. B. E. Kim and D. Y. M. Leung, "Significance of Skin Barrier Dysfunction in Atopic Dermatitis," *Allergy, Asthma & Immunology Research* 10, no. 3 (2018): 207–215.
5. Z. Celebi Sözen, L. Cevhertas, K. Nadeau, M. Akdis, and C. A. Akdis, "Environmental Factors in Epithelial Barrier Dysfunction," *Journal of Allergy and Clinical Immunology* 145, no. 6 (2020): 1517–1528.
6. S. Shimura, T. Takai, H. Iida, et al., "Epicutaneous Allergic Sensitization by Cooperation Between Allergen Protease Activity and Mechanical Skin Barrier Damage in Mice," *Journal of Investigative Dermatology* 136, no. 7 (2016): 1408–1417.
7. G. Imokawa, S. Akasaki, O. Kuno, et al., "Functions of Lipids on Human Skin," *Journal of Dispersion Science and Technology* 10, no. 4–5 (1989): 617–641.
8. G. Imokawa, S. Akasaki, M. Hattori, and N. Yoshizuka, "Selective Recovery of Deranged Water-Holding Properties by Stratum Corneum Lipids," *Journal of Investigative Dermatology* 87, no. 6 (1986): 758–761.
9. Y. Uchida, W. M. Holleran, and P. M. Elias, "On the Effects of Topical Synthetic Pseudoceramides: Comparison of Possible Keratinocyte Toxicities Provoked by the Pseudoceramides, PC104 and BIO391, and Natural Ceramides," *Journal of Dermatological Science* 51, no. 1 (2008): 37–43.
10. J. Verdrager, "Risk of Transmission of BSE via Drugs of Bovine Origin," *Lancet* 354, no. 9186 (1999): 1304–1305.
11. F. Le Joubioux, N. Bridiau, M. Sanekli, M. Graber, and T. Maugard, "Continuous Lipase-Catalyzed Production of Pseudo-Ceramides in a Packed-Bed Bioreactor," *Journal of Molecular Catalysis B: Enzymatic* 109 (2014): 143–153.
12. S. Murakami, T. Shimamoto, H. Nagano, et al., "Producing Human Ceramide-NS by Metabolic Engineering Using Yeast *Saccharomyces cerevisiae*," *Scientific Reports* 5 (2015): 16319.
13. C. Pérez-Rivero and J. P. López-Gómez, "Unlocking the Potential of Fermentation in Cosmetics: A Review," *Fermentation* 9, no. 5 (2023): 463.
14. M. Fujii, "The Pathogenic and Therapeutic Implications of Ceramide Abnormalities in Atopic Dermatitis," *Cells* 10, no. 9 (2021): 2386.
15. Y. Takagi, "Efficacy of Topical Application of a Skin Moisturizer Containing Pseudo-Ceramide and a Eucalyptus Leaf Extract on Atopic Dermatitis: A Review," *Journal of Clinical Medicine* 13, no. 6 (2024): 1749.
16. G. Imokawa, S. Akasaki, A. Kawamata, S. Yano, and N. Takaishi, "Water-Retaining Function in the Stratum Corneum and Its Recovery Properties by Synthetic Pseudoceramides," *Journal of the Society of Cosmetic Chemists* 40 (1989): 273–285.
17. H. Matsuki, K. Kiyokane, T. Matsuki, S. Sato, and G. Imokawa, "Re-evaluation of the Importance of Barrier Dysfunction in the Nonlesional Dry Skin of Atopic Dermatitis Patients Through the Use of Two Barrier Creams," *Exogenous Dermatology* 3, no. 6 (2006): 293–302.
18. K. Okoshi, Y. Kinugasa, S. Ito, et al., "Efficacy of Pseudo-Ceramide-Containing Steroid Lamellar Cream in Patients With Mild to Moderate Atopic Dermatitis: A Randomized, Double-Blind Study," *Dermatology and Therapy* 12, no. 8 (2022): 1823–1834.
19. M. Matsuoka, K. Okoshi, S. Ito, et al., "Efficacy of Heparinoid Cream Containing Pseudo-Ceramide for Remission of Atopic Dermatitis," *Clinical, Cosmetic and Investigational Dermatology* 14 (2021): 1839–1847.
20. M. Orita, M. Uchiyama, T. Hanamoto, et al., "Formation of Pseudo-Intercellular Lipids Membrane on the Skin Surface by the Alpha-Gel Holding a Large Amount of Water," *Journal of the Society of Cosmetic Chemists of Japan* 46, no. 1 (2012): 25–32.
21. S. X. Ge, E. W. Son, and R. Yao, "iDEP: An Integrated Web Application for Differential Expression and Pathway Analysis of RNA-Seq Data," *BMC Bioinformatics* 19, no. 1 (2018): 534.
22. H. Suto, H. Matsuda, K. Mitsuishi, et al., "NC/Nga Mice: A Mouse Model for Atopic Dermatitis," *International Archives of Allergy and Immunology* 120, no. Suppl 1 (1999): 70–75.
23. N. Takano, I. Arai, and M. Kurachi, "A Method to Induce Stable Atopic Dermatitis-Like Symptoms in NC/Nga Mice Housed With Skin-Lesioned Mice," *British Journal of Dermatology* 154, no. 3 (2006): 426–430.
24. M. Shiratori-Hayashi, K. Koga, H. Tozaki-Saitoh, et al., "STAT3-Dependent Reactive Astroglia in the Spinal Dorsal Horn Underlies Chronic Itch," *Nature Medicine* 21, no. 8 (2015): 927–931.
25. V. Singh and U. J. Phukan, "Interaction of Host and *Staphylococcus aureus* Protease-System Regulates Virulence and Pathogenicity," *Medical Microbiology and Immunology* 208, no. 5 (2019): 585–607.
26. T. Nakatsuji, T. H. Chen, A. M. Two, et al., "*Staphylococcus aureus* Exploits Epidermal Barrier Defects in Atopic Dermatitis to Trigger Cytokine Expression," *Journal of Investigative Dermatology* 136, no. 11 (2016): 2192–2200.
27. S. E. Poh, W. L. C. Koh, S. Y. D. Lim, et al., "Expression of *Staphylococcus aureus* Virulence Factors in Atopic Dermatitis," *JID Innovations* 2, no. 4 (2022): 100130.
28. Z. C. Yuan, W. D. Xu, X. Y. Liu, X. Y. Liu, A. F. Huang, and L. C. Su, "Biology of IL-36 Signaling and Its Role in Systemic Inflammatory Diseases," *Frontiers in Immunology* 10 (2019): 2532.
29. V. Todorović, Z. Su, C. B. Putman, et al., "Small Molecule IL-36γ Antagonist as a Novel Therapeutic Approach for Plaque Psoriasis," *Scientific Reports* 9, no. 1 (2019): 9089.
30. K. Shibata, H. Nomiyama, O. Yoshie, and S. Tanase, "Genome Diversification Mechanism of Rodent and Lagomorpha Chemokine Genes," *BioMed Research International* 2013 (2013): 856265.
31. Y. Hirasawa, T. Takai, T. Nakamura, et al., "*Staphylococcus aureus* Extracellular Protease Causes Epidermal Barrier Dysfunction," *Journal of Investigative Dermatology* 130, no. 2 (2010): 614–617.
32. T. Miyamoto, H. Nojima, T. Shinkado, T. Nakahashi, and Y. Kurai-shi, "Itch-Associated Response Induced by Experimental Dry Skin in Mice," *Japanese Journal of Pharmacology* 88, no. 3 (2002): 285–292.
33. L. S. Wong, Y. T. Yen, and C. H. Lee, "The Implications of Pruritogens in the Pathogenesis of Atopic Dermatitis," *International Journal of Molecular Sciences* 22, no. 13 (2021): 7227.
34. T. Buhl, A. Ikoma, C. Kempkes, et al., "Protease-Activated Receptor-2 Regulates Neuro-Epidermal Communication in Atopic Dermatitis," *Frontiers in Immunology* 11 (2020): 1740.
35. S. R. Wilson, L. Thé, L. M. Batia, et al., "The Epithelial Cell-Derived Atopic Dermatitis Cytokine TSLP Activates Neurons to Induce Itch," *Cell* 155, no. 2 (2013): 285–295.
36. T. Schmitt and R. H. H. Neubert, "State of the Art in Stratum Corneum Research: The Biophysical Properties of Ceramides," *Chemistry and Physics of Lipids* 216 (2018): 91–103.
37. M. Elias, S. Zhao, H. T. Le, et al., "IL-36 in Chronic Inflammation and Fibrosis—Bridging the Gap?," *Journal of Clinical Investigation* 131, no. 2 (2021): e144336.
38. A. M. D'Erme, D. Wilschmann-Theis, J. Wagenpfeil, et al., "IL-36γ (IL-1F9) is a Biomarker for Psoriasis Skin Lesions," *Journal of Investigative Dermatology* 135, no. 4 (2015): 1025–1032.
39. B. Wang, B. J. McHugh, A. Qureshi, et al., "IL-1β-Induced Protection of Keratinocytes Against *Staphylococcus aureus*-Secreted Proteases

Is Mediated by Human β -Defensin 2,” *Journal of Investigative Dermatology* 137, no. 1 (2017): 95–105.

40. K. Saito, Y. Iwata, H. Fukushima, et al., “IL-36 Receptor Antagonist Deficiency Resulted in Delayed Wound Healing due to Excessive Recruitment of Immune Cells,” *Scientific Reports* 10, no. 1 (2020): 14772.

41. S. Shukla, W. K. Cho, E. Elbasiony, R. B. Singh, S. K. Mittal, and S. K. Chauhan, “Non-Immune and Immune Functions of Interleukin-36 γ Suppress Epithelial Repair at the Ocular Surface,” *Federation of American Societies for Experimental Biology Journal* 36, no. 8 (2022): 202200174RR, <https://doi.org/10.1096/fj.202200174RR>.

42. K. Okoshi, S. Ito, M. Matsuoka, et al., “Combination of a Topical Anti-Inflammatory Drug and a Moisturizer, Both With a Lamellar Structure Containing Synthetic Pseudo-Ceramides, for the Treatment of Patients With Mild-To-Moderate Atopic Dermatitis,” *Clinical and Cosmetic Investigative Dermatology* 17 (2024): 1569–1578.

43. M. Tominaga and K. Takamori, “Peripheral Itch Sensitization in Atopic Dermatitis,” *Allergology International* 71, no. 3 (2022): 265–277.

44. L. F. Eichenfield, W. L. Tom, T. G. Berger, et al., “Guidelines of Care for the Management of Atopic Dermatitis: Section 2. Management and Treatment of Atopic Dermatitis With Topical Therapies,” *Journal of the American Academy of Dermatology* 71, no. 1 (2014): 116–132.

45. G. Imokawa, A. Abe, K. Jin, Y. Higaki, M. Kawashima, and A. Hidano, “Decreased Level of Ceramides in Stratum Corneum of Atopic Dermatitis: An Etiologic Factor in Atopic Dry Skin?,” *Journal of Investigative Dermatology* 96, no. 4 (1991): 523–526.

46. M. Henahan and A. De Benedetto, “Update on Protease-Activated Receptor 2 in Cutaneous Barrier, Differentiation, Tumorigenesis and Pigmentation, and Its Role in Related Dermatologic Diseases,” *Experimental Dermatology* 28, no. 8 (2019): 877–885.

47. T. Andoh, K. Tsujii, and Y. Kuraishi, “Increase in Pruritogenic Kallikrein 5 in the Skin of NC Mice With Chronic Dermatitis,” *Experimental Dermatology* 24, no. 12 (2015): 978–980.

48. C. J. Guo, M. R. Mack, L. K. Oetjen, et al., “Kallikrein 7 Promotes Atopic Dermatitis-Associated Itch Independently of Skin Inflammation,” *Journal of Investigative Dermatology* 140, no. 6 (2020): 1244–1252.e4.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.